

Table 4A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3 / 13 / 09

| Comparison of Racial and Ethnic Distribution by Clinic | | | | | | | | |
|---|--|--------------------------------------|--------------------------------------|-----------------------|---|---|-----------------------------|--------------------------------------|
| Clinic | Number Included in Analysis (3) | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaskan Native % | Asian/Pacific Islander % | Multiple Races % | All Other / Unknown % |
| 16Th St Co | 7,023 | 4.0 | 5.5 | 84.9 | 0.1 | 4.3 | 1.1 | 0.0 |
| Adams Cty | 586 | 84.3 | 0.3 | 9.2 | 0.0 | 0.0 | 6.1 | 0.0 |
| Ashland Cty | 545 | 79.3 | 0.0 | 2.4 | 6.1 | 0.2 | 12.1 | 0.0 |
| Aurora | 4,153 | 4.4 | 80.5 | 5.8 | 0.5 | 7.5 | 1.3 | 0.0 |
| Barron Cty | 1,145 | 84.6 | 4.8 | 8.4 | 0.5 | 0.2 | 1.5 | 0.0 |
| Bayfield Cty | 248 | 83.5 | 0.0 | 2.8 | 2.8 | 0.4 | 10.5 | 0.0 |
| Beloit | 2,060 | 46.7 | 15.2 | 34.9 | 0.0 | 0.4 | 2.7 | 0.0 |
| Buffalo Cty | 266 | 88.0 | 0.0 | 7.9 | 0.4 | 0.0 | 3.8 | 0.0 |
| Burnett Cty | 386 | 85.2 | 0.8 | 4.4 | 1.0 | 0.0 | 8.5 | 0.0 |
| Calumet Ct | 645 | 64.5 | 4.2 | 20.6 | 0.2 | 8.1 | 2.5 | 0.0 |
| Chippewa C | 1,419 | 90.6 | 0.7 | 2.6 | 0.6 | 1.1 | 4.4 | 0.1 |
| Clark Cty | 804 | 72.0 | 0.4 | 25.6 | 0.1 | 0.1 | 1.7 | 0.0 |
| Columbia C | 855 | 77.4 | 0.8 | 17.0 | 0.1 | 0.2 | 4.4 | 0.0 |
| Crawford Cty | 309 | 91.9 | 0.3 | 5.2 | 0.3 | 0.0 | 2.3 | 0.0 |
| Cudahy | 1,594 | 64.4 | 3.4 | 24.0 | 0.2 | 1.0 | 7.0 | 0.0 |
| Dane Cty | 6 | * | * | * | * | * | * | * |
| Dodge Cty | 1,361 | 65.3 | 1.2 | 28.7 | 0.1 | 0.4 | 4.2 | 0.0 |
| Door Cty | 479 | 81.8 | 2.3 | 13.4 | 0.4 | 0.2 | 1.9 | 0.0 |
| Douglas Cty | 1,233 | 80.9 | 0.6 | 4.6 | 1.1 | 0.8 | 11.9 | 0.0 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

| Comparison of Racial and Ethnic Distribution by Clinic | | | | | | | | |
|---|--|--------------------------------------|--------------------------------------|-----------------------|---|---|-----------------------------|--------------------------------------|
| Clinic | Number Included in Analysis (3) | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaskan Native % | Asian/Pacific Islander % | Multiple Races % | All Other / Unknown % |
| Dunn Cty | 927 | 78.4 | 0.6 | 5.9 | 0.1 | 9.9 | 5.0 | 0.0 |
| Eau Claire | 2,394 | 67.7 | 1.3 | 8.6 | 0.5 | 11.1 | 10.9 | 0.0 |
| Florence C | 128 | 90.6 | 0.0 | 5.5 | 1.6 | 0.0 | 2.3 | 0.0 |
| Fond Du La | 1,988 | 62.3 | 3.1 | 24.4 | 0.0 | 4.2 | 6.0 | 0.0 |
| Forest Cty | 225 | 90.2 | 0.0 | 2.2 | 2.2 | 0.0 | 5.3 | 0.0 |
| GLITC-Bad | 118 | 0.0 | 0.0 | 0.8 | 97.5 | 0.0 | 1.7 | 0.0 |
| GLITC-Bara | 56 | * | * | * | * | * | * | * |
| GLITC-BRF | 112 | 2.7 | 0.0 | 5.4 | 91.1 | 0.0 | 0.9 | 0.0 |
| GLITC-CEX | 33 | * | * | * | * | * | * | * |
| GLITC-LCO | 181 | 0.0 | 0.0 | 1.7 | 95.0 | 1.1 | 2.2 | 0.0 |
| GLITC-Mole | 39 | * | * | * | * | * | * | * |
| GLITC-Pete | 289 | 3.1 | 0.3 | 0.7 | 83.7 | 0.0 | 11.8 | 0.3 |
| GLITC-Pota | 82 | * | * | * | * | * | * | * |
| GLITC-Red | 115 | 6.1 | 0.9 | 4.3 | 87.0 | 0.0 | 1.7 | 0.0 |
| GLITC-St.C | 158 | 10.1 | 0.0 | 3.2 | 57.0 | 0.0 | 29.7 | 0.0 |
| GLITC-Stoc | 137 | 16.8 | 0.0 | 16.8 | 56.9 | 0.0 | 9.5 | 0.0 |
| Grant Cty | 1,243 | 92.1 | 0.8 | 5.1 | 0.0 | 0.2 | 1.7 | 0.0 |
| Green Cty | 655 | 80.2 | 0.2 | 17.3 | 0.0 | 0.3 | 2.1 | 0.0 |
| Green Lake | 315 | 71.7 | 0.3 | 21.3 | 0.0 | 2.2 | 4.1 | 0.3 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 3
Run Date: 3 / 13 / 09

| Comparison of Racial and Ethnic Distribution by Clinic | | | | | | | | |
|---|--|--------------------------------------|--------------------------------------|-----------------------|---|---|-----------------------------|--------------------------------------|
| Clinic | Number Included in Analysis (3) | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaskan Native % | Asian/Pacific Islander % | Multiple Races % | All Other / Unknown % |
| Greenfield | 886 | 57.8 | 3.6 | 27.5 | 0.3 | 1.8 | 8.9 | 0.0 |
| Hlth & Nut | 3,762 | 30.4 | 20.3 | 40.0 | 0.1 | 0.3 | 8.9 | 0.0 |
| Iowa Cty | 356 | 85.1 | 1.4 | 8.1 | 0.3 | 0.6 | 4.5 | 0.0 |
| Iron Cty | 97 | * | * | * | * | * | * | * |
| Jackson Ct | 570 | 81.1 | 0.7 | 6.1 | 8.2 | 0.9 | 2.8 | 0.2 |
| Janesville | 2,094 | 78.0 | 4.2 | 13.1 | 0.1 | 1.3 | 3.3 | 0.0 |
| Jefferson | 1,725 | 51.2 | 0.7 | 45.6 | 0.1 | 0.3 | 2.0 | 0.1 |
| Juneau | 725 | 87.3 | 0.6 | 6.9 | 0.1 | 0.7 | 4.4 | 0.0 |
| Kenosha | 4,128 | 34.8 | 17.5 | 40.5 | 0.4 | 0.6 | 6.2 | 0.0 |
| Kewaunee C | 323 | 71.2 | 1.5 | 23.8 | 0.3 | 0.0 | 3.1 | 0.0 |
| La Crosse | 2,425 | 67.3 | 3.5 | 4.2 | 0.4 | 14.9 | 9.7 | 0.0 |
| Langlade | 488 | 87.1 | 0.6 | 8.2 | 0.0 | 0.0 | 4.1 | 0.0 |
| Lincoln | 475 | 86.9 | 0.2 | 7.2 | 0.4 | 0.0 | 5.3 | 0.0 |
| Madison WI | 7,344 | 27.4 | 21.5 | 36.0 | 0.2 | 7.1 | 7.8 | 0.0 |
| Manitowoc | 1,660 | 58.1 | 1.3 | 23.0 | 0.7 | 10.6 | 6.2 | 0.0 |
| Marathon | 2,683 | 55.7 | 1.8 | 9.9 | 0.3 | 25.2 | 7.1 | 0.0 |
| Marinette | 1,051 | 94.9 | 0.1 | 2.5 | 0.5 | 0.1 | 2.0 | 0.0 |
| Marquette | 254 | 74.8 | 0.4 | 20.5 | 0.4 | 0.0 | 3.9 | 0.0 |
| Menominee | 588 | 0.2 | 0.0 | 15.8 | 34.2 | 0.2 | 49.3 | 0.3 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 4
Run Date: 3 / 13 / 09

| Comparison of Racial and Ethnic Distribution by Clinic | | | | | | | | |
|---|--|--------------------------------------|--------------------------------------|-----------------------|---|---|-----------------------------|--------------------------------------|
| Clinic | Number Included in Analysis (3) | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaskan Native % | Asian/Pacific Islander % | Multiple Races % | All Other / Unknown % |
| MilwaukeeJ | 2,591 | 14.2 | 7.6 | 70.7 | 0.8 | 3.0 | 3.6 | 0.0 |
| MilwaukeeK | 2,338 | 1.0 | 93.1 | 2.8 | 0.1 | 0.9 | 2.1 | 0.1 |
| MilwaukeeN | 3,663 | 8.5 | 70.6 | 7.3 | 0.1 | 6.5 | 7.0 | 0.1 |
| MilwaukMLK | 2,967 | 3.1 | 83.0 | 7.9 | 0.2 | 0.7 | 4.9 | 0.1 |
| Monroe Cou | 1,153 | 75.0 | 1.1 | 19.1 | 0.6 | 1.1 | 3.0 | 0.0 |
| N.E.W. East | 3,058 | 32.7 | 4.4 | 54.4 | 1.5 | 4.1 | 2.9 | 0.0 |
| N.E.W. West | 3,026 | 56.6 | 6.6 | 17.5 | 2.7 | 10.2 | 6.3 | 0.0 |
| Neenah | 1,273 | 63.4 | 3.2 | 24.7 | 0.2 | 3.6 | 4.9 | 0.0 |
| Oconto Cty | 631 | 91.6 | 0.2 | 5.7 | 0.3 | 0.2 | 2.1 | 0.0 |
| Oneida Cty | 633 | 91.0 | 0.3 | 4.4 | 0.6 | 0.3 | 3.3 | 0.0 |
| Oneida Tri | 579 | 8.3 | 0.5 | 22.8 | 54.4 | 0.7 | 13.3 | 0.0 |
| Oshkosh | 1,340 | 63.0 | 4.4 | 10.7 | 0.4 | 12.4 | 9.0 | 0.0 |
| Outagamie | 2,593 | 56.8 | 2.4 | 21.4 | 0.7 | 13.3 | 5.3 | 0.0 |
| Ozaukee | 498 | 75.1 | 3.0 | 12.2 | 0.4 | 0.6 | 8.6 | 0.0 |
| Pepin | 152 | 87.5 | 0.7 | 6.6 | 0.7 | 0.7 | 3.9 | 0.0 |
| Pierce Cty | 724 | 86.7 | 1.1 | 5.2 | 0.6 | 0.6 | 5.8 | 0.0 |
| Polk Cty | 982 | 92.4 | 0.2 | 3.5 | 0.6 | 0.1 | 3.3 | 0.0 |
| Portage Ct | 1,210 | 66.6 | 0.6 | 15.7 | 0.2 | 13.6 | 3.3 | 0.0 |
| Price Cty | 475 | 89.5 | 0.4 | 3.4 | 0.2 | 3.4 | 3.2 | 0.0 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 5
Run Date: 3 / 13 / 09

| Comparison of Racial and Ethnic Distribution by Clinic | | | | | | | | |
|---|--|--------------------------------------|--------------------------------------|-----------------------|---|---|-----------------------------|--------------------------------------|
| Clinic | Number Included in Analysis (3) | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaskan Native % | Asian/Pacific Islander % | Multiple Races % | All Other / Unknown % |
| Racine Hea | 1,806 | 15.5 | 38.4 | 38.4 | 0.2 | 0.3 | 6.5 | 0.8 |
| Richland | 398 | 87.7 | 0.0 | 10.8 | 0.3 | 0.3 | 1.0 | 0.0 |
| Rusk Cty | 414 | 89.4 | 1.2 | 4.3 | 0.0 | 0.0 | 5.1 | 0.0 |
| Sauk Cty | 1,403 | 76.3 | 1.1 | 19.8 | 0.9 | 0.0 | 1.9 | 0.0 |
| Sawyer Cty | 460 | 70.0 | 1.5 | 5.0 | 19.6 | 0.7 | 3.3 | 0.0 |
| Seeds Of H | 3,282 | 14.7 | 3.6 | 72.8 | 0.4 | 1.4 | 7.1 | 0.0 |
| Shawano Ct | 834 | 74.8 | 0.6 | 9.7 | 6.5 | 0.4 | 8.0 | 0.0 |
| Sheboygan | 2,377 | 45.8 | 2.7 | 29.7 | 0.0 | 17.7 | 4.0 | 0.0 |
| St. Croix | 1,196 | 82.2 | 1.3 | 8.5 | 0.3 | 2.5 | 5.3 | 0.0 |
| SWCAP | 220 | 75.9 | 1.4 | 20.9 | 0.0 | 0.0 | 1.8 | 0.0 |
| Taylor Cty | 453 | 85.2 | 0.2 | 10.6 | 0.0 | 0.0 | 4.0 | 0.0 |
| Trempealeau | 800 | 62.1 | 0.1 | 35.3 | 0.4 | 0.3 | 1.9 | 0.0 |
| Vernon Cou | 758 | 88.0 | 0.7 | 8.0 | 0.1 | 0.4 | 2.8 | 0.0 |
| Vilas Cty | 280 | 92.5 | 0.7 | 3.6 | 1.8 | 0.0 | 1.4 | 0.0 |
| Walworth C | 2,062 | 45.1 | 1.1 | 49.0 | 0.6 | 0.8 | 3.3 | 0.0 |
| Washburn C | 449 | 87.8 | 0.0 | 5.6 | 1.6 | 1.1 | 4.0 | 0.0 |
| Washington | 1,522 | 75.1 | 3.0 | 15.0 | 0.3 | 0.7 | 5.8 | 0.0 |
| Waukesha C | 2,439 | 48.8 | 5.7 | 36.9 | 0.4 | 1.2 | 7.1 | 0.0 |
| Waupaca Ct | 853 | 85.7 | 0.0 | 12.7 | 0.0 | 0.1 | 1.5 | 0.0 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 6
Run Date: 3 / 13 / 09

| Comparison of Racial and Ethnic Distribution by Clinic | | | | | | | | |
|---|--|--------------------------------------|--------------------------------------|-----------------------|---|---|-----------------------------|--------------------------------------|
| Clinic | Number Included in Analysis (3) | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaskan Native % | Asian/Pacific Islander % | Multiple Races % | All Other / Unknown % |
| Waushara | 501 | 54.9 | 0.4 | 42.5 | 0.2 | 0.2 | 1.8 | 0.0 |
| WC Capitol | 3,873 | 6.9 | 81.1 | 3.6 | 0.1 | 5.0 | 3.2 | 0.1 |
| WC North | 2,185 | 3.8 | 79.5 | 3.6 | 0.0 | 10.0 | 3.0 | 0.0 |
| WC Teutonia | 2,374 | 1.1 | 92.6 | 2.7 | 0.0 | 0.5 | 2.9 | 0.2 |
| West Allis | 1,341 | 54.1 | 7.8 | 24.1 | 0.5 | 1.7 | 11.8 | 0.0 |
| Wood Cty | 1,536 | 81.3 | 0.7 | 9.3 | 0.3 | 4.1 | 4.4 | 0.0 |
| Wisconsin | 129,648 | 45.1 | 18.5 | 24.9 | 1.7 | 4.4 | 5.2 | 0.0 |
| Nation (Prior Year) | 7,995,600 | 31.6 | 19.4 | 40.9 | 1.0 | 2.5 | 1.8 | 2.7 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3 / 13 / 09

| Comparison of Age Distribution by Clinic | | | | |
|---|--|---------------------------------|----------------------------------|----------------------------------|
| Clinic | Number Included in Analysis (3,4) | 0 - 11 Months % | 12 - 23 Months % | 24 - 59 Months % |
| 16Th St Co | 7,023 | 30.5 | 19.8 | 49.7 |
| Adams Cty | 586 | 34.6 | 20.0 | 45.4 |
| Ashland Cty | 545 | 29.0 | 22.6 | 48.4 |
| Aurora | 4,153 | 48.7 | 14.9 | 36.4 |
| Barron Cty | 1,145 | 36.6 | 21.5 | 41.9 |
| Bayfield Cty | 248 | 31.0 | 19.4 | 49.6 |
| Beloit | 2,060 | 36.1 | 22.0 | 41.8 |
| Buffalo Cty | 266 | 37.6 | 17.7 | 44.7 |
| Burnett Cty | 386 | 31.9 | 23.6 | 44.6 |
| Calumet Ct | 645 | 30.5 | 22.3 | 47.1 |
| Chippewa C | 1,419 | 31.1 | 22.3 | 46.6 |
| Clark Cty | 804 | 33.6 | 20.3 | 46.1 |
| Columbia C | 855 | 37.3 | 21.8 | 40.9 |
| Crawford Cty | 309 | 37.5 | 19.7 | 42.7 |
| Cudahy | 1,594 | 34.4 | 25.8 | 39.8 |
| Dane Cty | 6 | * | * | * |
| Dodge Cty | 1,361 | 34.3 | 20.6 | 45.1 |
| Door Cty | 479 | 37.0 | 15.4 | 47.6 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 2
Run Date: 3 / 13 / 09

| Comparison of Age Distribution by Clinic | | | | |
|---|--|---------------------------------|----------------------------------|----------------------------------|
| Clinic | Number Included in Analysis (3,4) | 0 - 11 Months % | 12 - 23 Months % | 24 - 59 Months % |
| Douglas Cty | 1,233 | 36.2 | 20.9 | 42.9 |
| Dunn Cty | 927 | 34.0 | 19.1 | 46.9 |
| Eau Claire | 2,394 | 35.3 | 19.5 | 45.3 |
| Florence C | 128 | 32.0 | 20.3 | 47.7 |
| Fond Du La | 1,988 | 32.2 | 22.4 | 45.4 |
| Forest Cty | 225 | 33.3 | 20.0 | 46.7 |
| GLITC-Bad | 118 | 26.3 | 22.9 | 50.8 |
| GLITC-Bara | 56 | * | * | * |
| GLITC-BRF | 112 | 21.4 | 28.6 | 50.0 |
| GLITC-CEX | 33 | * | * | * |
| GLITC-LCO | 181 | 26.5 | 26.0 | 47.5 |
| GLITC-Mole | 39 | * | * | * |
| GLITC-Pete | 289 | 28.7 | 23.2 | 48.1 |
| GLITC-Pota | 82 | * | * | * |
| GLITC-Red | 115 | 29.6 | 21.7 | 48.7 |
| GLITC-St.C | 158 | 27.2 | 19.0 | 53.8 |
| GLITC-Stoc | 137 | 32.8 | 15.3 | 51.8 |
| Grant Cty | 1,243 | 33.7 | 20.1 | 46.2 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 3
Run Date: 3 / 13 / 09

| Comparison of Age Distribution by Clinic | | | | |
|---|--|---------------------------------|----------------------------------|----------------------------------|
| Clinic | Number Included in Analysis (3,4) | 0 - 11 Months % | 12 - 23 Months % | 24 - 59 Months % |
| Green Cty | 655 | 33.4 | 21.4 | 45.2 |
| Green Lake | 315 | 39.0 | 15.9 | 45.1 |
| Greenfield | 886 | 38.1 | 23.3 | 38.6 |
| Hlth & Nut | 3,762 | 39.5 | 19.2 | 41.3 |
| Iowa Cty | 356 | 42.7 | 20.5 | 36.8 |
| Iron Cty | 97 | * | * | * |
| Jackson Ct | 570 | 32.3 | 18.6 | 49.1 |
| Janesville | 2,094 | 40.9 | 17.9 | 41.2 |
| Jefferson | 1,725 | 35.9 | 19.7 | 44.3 |
| Juneau | 725 | 34.9 | 21.0 | 44.1 |
| Kenosha | 4,128 | 37.3 | 22.1 | 40.6 |
| Kewaunee C | 323 | 39.6 | 15.8 | 44.6 |
| La Crosse | 2,425 | 34.0 | 22.1 | 44.0 |
| Langlade | 488 | 33.4 | 20.5 | 46.1 |
| Lincoln | 475 | 33.9 | 20.6 | 45.5 |
| Madison WI | 7,344 | 36.1 | 21.8 | 42.1 |
| Manitowoc | 1,660 | 36.9 | 18.4 | 44.7 |
| Marathon | 2,683 | 34.2 | 21.7 | 44.1 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 4
Run Date: 3 / 13 / 09

| Comparison of Age Distribution by Clinic | | | | |
|---|--|---------------------------------|----------------------------------|----------------------------------|
| Clinic | Number Included in Analysis (3,4) | 0 - 11 Months % | 12 - 23 Months % | 24 - 59 Months % |
| Marinette | 1,051 | 31.8 | 22.5 | 45.8 |
| Marquette | 254 | 37.4 | 16.5 | 46.1 |
| Menominee | 588 | 31.5 | 22.3 | 46.3 |
| MilwaukeeJ | 2,591 | 29.5 | 22.5 | 48.0 |
| MilwaukeeK | 2,338 | 27.6 | 24.9 | 47.4 |
| MilwaukeeN | 3,663 | 32.6 | 25.1 | 42.3 |
| MilwaukMLK | 2,967 | 30.5 | 25.3 | 44.2 |
| Monroe Cou | 1,153 | 36.2 | 21.4 | 42.4 |
| N.E.W. East | 3,058 | 29.0 | 23.4 | 47.5 |
| N.E.W. West | 3,026 | 34.6 | 21.9 | 43.5 |
| Neenah | 1,273 | 40.1 | 20.2 | 39.7 |
| Oconto Cty | 631 | 35.3 | 18.5 | 46.1 |
| Oneida Cty | 633 | 41.1 | 16.7 | 42.2 |
| Oneida Tri | 579 | 33.7 | 25.6 | 40.8 |
| Oshkosh | 1,340 | 38.1 | 20.1 | 41.8 |
| Outagamie | 2,593 | 32.5 | 22.0 | 45.5 |
| Ozaukee | 498 | 36.1 | 22.5 | 41.4 |
| Pepin | 152 | 32.2 | 13.8 | 53.9 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 5
Run Date: 3 / 13 / 09

| Comparison of Age Distribution by Clinic | | | | |
|---|--|---------------------------------|----------------------------------|----------------------------------|
| Clinic | Number Included in Analysis (3,4) | 0 - 11 Months % | 12 - 23 Months % | 24 - 59 Months % |
| Pierce Cty | 724 | 32.6 | 20.4 | 47.0 |
| Polk Cty | 982 | 35.0 | 23.4 | 41.5 |
| Portage Ct | 1,210 | 38.0 | 19.8 | 42.2 |
| Price Cty | 475 | 29.3 | 19.8 | 50.9 |
| Racine Hea | 1,806 | 35.3 | 19.7 | 45.1 |
| Richland | 398 | 36.2 | 20.1 | 43.7 |
| Rusk Cty | 414 | 34.8 | 19.3 | 45.9 |
| Sauk Cty | 1,403 | 36.6 | 20.1 | 43.3 |
| Sawyer Cty | 460 | 34.6 | 18.0 | 47.4 |
| Seeds Of H | 3,282 | 30.7 | 25.2 | 44.1 |
| Shawano Ct | 834 | 28.5 | 26.5 | 45.0 |
| Sheboygan | 2,377 | 31.0 | 23.4 | 45.6 |
| St. Croix | 1,196 | 37.7 | 18.5 | 43.8 |
| SWCAP | 220 | 35.0 | 21.8 | 43.2 |
| Taylor Cty | 453 | 37.3 | 20.1 | 42.6 |
| Trempealeau | 800 | 36.3 | 20.6 | 43.1 |
| Vernon Cou | 758 | 35.2 | 21.1 | 43.7 |
| Vilas Cty | 280 | 36.4 | 21.1 | 42.5 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 6
Run Date: 3 / 13 / 09

| Comparison of Age Distribution by Clinic | | | | |
|---|--|---------------------------------|----------------------------------|----------------------------------|
| Clinic | Number Included in Analysis (3,4) | 0 - 11 Months % | 12 - 23 Months % | 24 - 59 Months % |
| Walworth C | 2,062 | 39.8 | 18.8 | 41.4 |
| Washburn C | 449 | 31.6 | 21.6 | 46.8 |
| Washington | 1,522 | 35.4 | 21.6 | 43.0 |
| Waukesha C | 2,439 | 41.2 | 18.4 | 40.3 |
| Waupaca Ct | 853 | 37.3 | 23.3 | 39.4 |
| Waushara | 501 | 37.9 | 17.4 | 44.7 |
| WC Capitol | 3,873 | 37.8 | 20.8 | 41.4 |
| WC North | 2,185 | 33.8 | 22.2 | 43.9 |
| WC Teutonia | 2,374 | 32.1 | 22.2 | 45.7 |
| West Allis | 1,341 | 38.6 | 22.7 | 38.6 |
| Wood Cty | 1,536 | 41.9 | 18.4 | 39.7 |
| Wisconsin | 129,648 | 35.0 | 21.2 | 43.8 |
| Nation (Prior Year) | 7,995,600 | 37.6 | 22.1 | 40.3 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | | | |
|--|-----------------|------------------------|------------------------|-------------------------|---------------------|-------------------------|----------------------|--------------------------------|----------------------|----------------------|-----------------------------|--------|-------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | | |
| | Low | | High | Short Stature (6) | | Under- weight (7) | Obese (7) | ≥ 2 Yrs Over- weight (7) | | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- (Rank) | % <95th (Rank) | % ≥95th (Rank) | Number | % (Rank) |
| 16Th St Co | 1,449 | 6.6 (18) | 7.5 (23) | 6,888 | 7.0 (86) | 1.1 (3) | 21.6 (89) | 3,433 | 21.9 (81) | 24.0 (84) | | 5,337 | 11.6 (66) |
| Adams Cty | 139 | 11.5 (64) | 14.4 (70) | 571 | 6.1 (69) | 2.6 (35) | 13.7 (56) | 264 | 17.8 (59) | 12.5 (35) | | 423 | 7.1 (33) |
| Ashland Cty | 111 | 3.6 (2) | 14.4 (70) | 540 | 5.4 (52) | 2.2 (19) | 11.7 (32) | 259 | 12.7 (13) | 10.8 (17) | | 424 | 5.7 (25) |
| Aurora | 1,543 | 9.5 (53) | 4.3 (4) | 4,083 | 4.4 (25) | 7.0 (94) | 9.3 (4) | 1,478 | 14.7 (22) | 10.5 (14) | | 2,191 | 28.5 (91) |
| Barron Cty | 279 | 8.2 (38) | 9.7 (52) | 1,125 | 4.6 (31) | 2.7 (38) | 12.3 (40) | 488 | 16.8 (46) | 13.9 (52) | | 814 | 9.7 (55) |
| Bayfield Cty | 55 | * * | * * | 245 | 2.0 (3) | 3.7 (64) | 11.0 (20) | 124 | 19.4 (72) | 12.9 (40) | | 187 | 8.6 (45) |
| Beloit | 498 | 6.4 (16) | 5.8 (10) | 2,015 | 4.8 (36) | 3.4 (57) | 16.3 (79) | 845 | 16.2 (38) | 16.7 (71) | | 1,449 | 8.1 (39) |
| Buffalo Cty | 67 | * * | * * | 261 | 2.3 (4) | 2.7 (38) | 10.7 (18) | 118 | 16.1 (37) | 13.6 (50) | | 190 | 12.1 (68) |
| Burnett Cty | 88 | * * | * * | 381 | 4.2 (20) | 3.7 (64) | 9.4 (6) | 168 | 14.9 (24) | 8.9 (7) | | 288 | 2.1 (1) |
| Calumet Ct | 148 | 8.8 (47) | 8.1 (32) | 633 | 6.5 (75) | 5.5 (88) | 12.5 (42) | 302 | 16.6 (43) | 15.9 (66) | | 482 | 4.6 (16) |
| Chippewa C | 335 | 6.6 (18) | 8.1 (32) | 1,401 | 6.9 (84) | 2.4 (28) | 12.8 (48) | 646 | 17.5 (52) | 13.3 (45) | | 1,052 | 12.5 (70) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 2
Run Date: 3 / 13 / 09

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | | |
|--|-----------------|------------------------|------------------------|-------------------------|---------------------|-------------------------|----------------------|--------------------------------|----------------------------|----------------------|-----------------------------|-------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | High | | Short Stature (6) | | Under- weight (7) | Obese (7) | ≥ 2 Yrs Over- weight (7) | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- <95th (Rank) | % ≥95th (Rank) | Number | % (Rank) |
| Clark Cty | 180 | 9.4 (52) | 9.4 (49) | 788 | 5.5 (57) | 2.3 (23) | 13.1 (50) | 373 | 21.7 (80) | 11.0 (18) | 582 | 13.9 (76) |
| Columbia C | 222 | 9.0 (50) | 9.9 (55) | 847 | 5.5 (57) | 4.3 (76) | 10.4 (14) | 342 | 16.4 (41) | 12.0 (28) | 598 | 8.9 (48) |
| Crawford Cty | 88 | * * | * * | 306 | 6.2 (71) | 3.9 (71) | 8.8 (2) | 133 | 12.0 (7) | 12.0 (28) | 213 | 11.3 (65) |
| Cudahy | 429 | 8.2 (38) | 9.1 (43) | 1,548 | 6.0 (65) | 3.2 (54) | 11.6 (31) | 629 | 15.4 (28) | 12.4 (32) | 1,105 | 9.3 (53) |
| Dane Cty | 0 | * * | * * | 4 | * * | * * | * * | 0 | * * | * * | 6 | * * |
| Dodge Cty | 334 | 8.4 (42) | 6.3 (14) | 1,329 | 4.7 (33) | 4.4 (80) | 9.9 (11) | 601 | 12.3 (9) | 7.8 (5) | 966 | 5.0 (19) |
| Door Cty | 124 | 7.3 (27) | 10.5 (59) | 474 | 4.0 (17) | 3.8 (67) | 8.9 (3) | 229 | 16.6 (43) | 13.5 (48) | 339 | 3.8 (11) |
| Douglas Cty | 302 | 4.6 (7) | 13.6 (67) | 1,218 | 3.9 (15) | 3.8 (67) | 14.0 (59) | 529 | 19.3 (70) | 17.0 (73) | 821 | 13.4 (73) |
| Dunn Cty | 202 | 5.4 (8) | 9.4 (49) | 907 | 6.0 (65) | 3.4 (57) | 10.9 (19) | 420 | 19.5 (73) | 10.7 (15) | 677 | 4.9 (17) |
| Eau Claire | 578 | 5.5 (10) | 9.7 (52) | 2,370 | 5.4 (52) | 2.5 (31) | 10.2 (12) | 1,070 | 17.5 (52) | 11.7 (23) | 1,724 | 11.8 (67) |
| Florence C | 26 | * * | * * | 125 | 8.8 (94) | 0.0 (1) | 11.2 (24) | 62 | * * | * * | 86 | * * |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 3
Run Date: 3 / 13 / 09

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | | | |
|--|-----------------|------------------------|------------------------|-------------------------|---------------------|-------------------------|----------------------|--------------------------------|----------------------|----------------------|-----------------------------|--------|-------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | | |
| | Low | | High | Short Stature (6) | | Under- weight (7) | Obese (7) | ≥ 2 Yrs Over- weight (7) | | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- (Rank) | % <95th (Rank) | % ≥95th (Rank) | Number | % (Rank) |
| Fond Du La | 455 | 9.9 (55) | 8.1 (32) | 1,946 | 7.0 (86) | 3.4 (57) | 10.5 (17) | 898 | 14.7 (22) | 9.7 (9) | | 1,467 | 8.9 (48) |
| Forest Cty | 47 | * * | * * | 218 | 5.0 (43) | 3.2 (54) | 16.1 (78) | 106 | 13.2 (15) | 19.8 (79) | | 169 | 10.1 (57) |
| GLITC-Bad | 22 | * * | * * | 115 | 1.7 (2) | 1.7 (10) | 18.3 (87) | 61 | * * | * * | | 95 | * * |
| GLITC-Bara | 11 | * * | * * | 53 | * * | * * | * * | 21 | * * | * * | | 39 | * * |
| GLITC-BRF | 18 | * * | * * | 106 | 5.7 (61) | 2.8 (43) | 23.6 (93) | 49 | * * | * * | | 92 | * * |
| GLITC-CEX | 6 | * * | * * | 32 | * * | * * | * * | 17 | * * | * * | | 25 | * * |
| GLITC-LCO | 35 | * * | * * | 178 | 2.8 (8) | 2.8 (43) | 16.3 (79) | 80 | * * | * * | | 135 | 15.6 (81) |
| GLITC-Mole | 6 | * * | * * | 39 | * * | * * | * * | 22 | * * | * * | | 32 | * * |
| GLITC-Pete | 50 | * * | * * | 273 | 3.7 (12) | 2.2 (19) | 24.9 (94) | 131 | 24.4 (86) | 20.6 (81) | | 220 | 3.2 (4) |
| GLITC-Pota | 16 | * * | * * | 77 | * * | * * | * * | 35 | * * | * * | | 47 | * * |
| GLITC-Red | 27 | * * | * * | 109 | 2.8 (8) | 6.4 (90) | 22.0 (91) | 51 | * * | * * | | 83 | * * |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | |
|---|-----------------|------------------------|------------------------|-------------------------|-------------------------|---------------------|----------------------|--------------------------------|----------------------------|-----------------------------|--------------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | High | | Short Stature (6) | Under- weight (7) | Obese (7) | | ≥ 2 Yrs Over- weight (7) | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- <95th (Rank) | % ≥95th (Rank) | Number % (Rank) |
| GLITC-St.C | 30 | * * | * * | 153 | 4.6 (31) | 1.3 (6) | 29.4 (95) | 79 | * * | * * | 122 8.2 (40) |
| GLITC-Stoc | 25 | * * | * * | 134 | 0.7 (1) | 0.0 (1) | 18.7 (88) | 73 | * * | * * | 107 16.8 (83) |
| Grant Cty | 286 | 7.0 (25) | 11.5 (61) | 1,223 | 4.5 (27) | 3.8 (67) | 11.0 (20) | 570 | 15.3 (27) | 12.1 (31) | 940 3.2 (4) |
| Green Cty | 148 | 10.1 (56) | 12.2 (63) | 642 | 7.9 (90) | 2.6 (35) | 18.1 (85) | 285 | 17.9 (61) | 24.9 (85) | 464 10.6 (62) |
| Green Lake | 69 | * * | * * | 307 | 2.6 (5) | 2.6 (35) | 13.0 (49) | 136 | 12.5 (11) | 14.0 (54) | 221 8.6 (45) |
| Greenfield | 271 | 8.5 (43) | 9.2 (45) | 862 | 6.7 (80) | 2.8 (43) | 11.4 (29) | 334 | 17.1 (48) | 12.0 (28) | 578 15.2 (80) |
| Hlth & Nut | 1,028 | 11.1 (63) | 6.9 (21) | 3,659 | 5.9 (63) | 3.6 (62) | 13.6 (54) | 1,533 | 16.4 (41) | 13.6 (50) | 2,613 8.2 (40) |
| Iowa Cty | 110 | 12.7 (70) | 6.4 (15) | 346 | 6.6 (77) | 1.7 (10) | 11.8 (34) | 131 | 17.6 (55) | 9.9 (11) | 232 8.6 (45) |
| Iron Cty | 26 | * * | * * | 94 | * * | * * | * * | 45 | * * | * * | 67 * * |
| Jackson Ct | 119 | 4.2 (4) | 14.3 (68) | 566 | 6.9 (84) | 1.8 (14) | 12.2 (37) | 280 | 20.7 (79) | 13.9 (52) | 390 7.7 (38) |
| Janesville | 569 | 7.9 (36) | 8.4 (37) | 2,059 | 3.8 (14) | 4.3 (76) | 11.9 (36) | 847 | 17.8 (59) | 12.5 (35) | 1,391 8.9 (48) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 5
Run Date: 3 / 13 / 09

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | |
|---|-----------------|------------------------|------------------------|-------------------------|-------------------------|---------------------|----------------------|--------------------------------|----------------------------|-----------------------------|--------------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | High | | Short Stature (6) | Under- weight (7) | Obese (7) | | ≥ 2 Yrs Over- weight (7) | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- <95th (Rank) | % ≥95th (Rank) | Number % (Rank) |
| Jefferson | 402 | 6.2 (13) | 8.5 (39) | 1,686 | 4.3 (22) | 2.9 (47) | 12.4 (41) | 764 | 17.5 (52) | 14.5 (58) | 1,236 5.7 (25) |
| Juneau | 155 | 5.8 (11) | 7.7 (25) | 701 | 5.1 (48) | 1.7 (10) | 15.7 (76) | 319 | 15.4 (28) | 16.3 (70) | 531 7.2 (34) |
| Kenosha | 1,107 | 10.7 (60) | 6.8 (19) | 4,028 | 7.9 (90) | 1.8 (14) | 15.5 (75) | 1,652 | 15.9 (35) | 12.5 (35) | 2,803 12.5 (70) |
| Kewaunee C | 77 | * * | * * | 321 | 5.3 (51) | 1.2 (5) | 18.1 (85) | 143 | 20.3 (77) | 13.3 (45) | 229 14.8 (79) |
| La Crosse | 600 | 8.3 (40) | 8.0 (29) | 2,363 | 6.7 (80) | 3.9 (71) | 11.3 (25) | 1,059 | 19.9 (76) | 11.0 (18) | 1,764 6.6 (31) |
| Langlade | 113 | 8.0 (37) | 8.0 (29) | 484 | 5.0 (43) | 2.3 (23) | 13.2 (52) | 225 | 20.4 (78) | 18.7 (77) | 370 19.2 (86) |
| Lincoln | 110 | 3.6 (2) | 9.1 (43) | 468 | 5.6 (60) | 1.7 (10) | 15.0 (71) | 216 | 16.7 (45) | 17.1 (74) | 356 5.1 (20) |
| Madison WI | 1,882 | 7.5 (29) | 8.4 (37) | 7,136 | 4.5 (27) | 3.1 (52) | 12.6 (44) | 3,064 | 12.2 (8) | 10.2 (13) | 5,126 11.0 (63) |
| Manitowoc | 403 | 10.7 (60) | 6.5 (16) | 1,641 | 4.8 (36) | 3.4 (57) | 14.0 (59) | 737 | 18.7 (67) | 15.6 (65) | 1,140 8.2 (40) |
| Marathon | 622 | 7.7 (33) | 5.1 (7) | 2,633 | 8.3 (93) | 1.5 (7) | 14.0 (59) | 1,167 | 18.3 (62) | 13.3 (45) | 2,003 4.1 (14) |
| Marinette | 222 | 5.4 (8) | 9.0 (42) | 1,030 | 4.7 (33) | 6.7 (92) | 8.6 (1) | 474 | 9.3 (2) | 5.3 (1) | 769 3.9 (13) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 6
Run Date: 3 / 13 / 09

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | | |
|--|-----------------|------------------------|------------------------|-------------------------|---------------------|-------------------------|----------------------|--------------------------------|----------------------------|----------------------|-----------------------------|-------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | | High | Short Stature (6) | | Under- weight (7) | Obese (7) | ≥ 2 Yrs Over- weight (7) | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- <95th (Rank) | % ≥95th (Rank) | Number | % (Rank) |
| Marquette | 56 | * * | * * | 249 | 6.8 (82) | 2.0 (17) | 17.7 (83) | 117 | 19.7 (75) | 15.4 (62) | 178 | 5.1 (20) |
| Menominee | 147 | 3.4 (1) | 14.3 (68) | 570 | 2.6 (5) | 1.1 (3) | 21.6 (89) | 263 | 23.2 (85) | 21.3 (83) | 424 | 5.9 (28) |
| MilwaukeeJ | 589 | 7.3 (27) | 8.3 (36) | 2,526 | 6.3 (73) | 3.4 (57) | 14.1 (62) | 1,220 | 16.2 (38) | 15.0 (60) | 1,940 | 13.7 (75) |
| MilwaukeeK | 498 | 12.0 (68) | 3.8 (3) | 2,288 | 6.2 (71) | 3.8 (67) | 11.5 (30) | 1,093 | 15.6 (31) | 11.4 (21) | 1,789 | 24.4 (89) |
| MilwaukeeN | 899 | 11.7 (65) | 4.7 (6) | 3,535 | 9.1 (95) | 5.3 (86) | 13.6 (54) | 1,517 | 16.3 (40) | 13.5 (48) | 2,625 | 23.2 (88) |
| MilwaukMLK | 629 | 11.8 (66) | 4.5 (5) | 2,892 | 4.7 (33) | 6.9 (93) | 9.8 (10) | 1,280 | 10.1 (4) | 6.4 (2) | 2,234 | 19.1 (85) |
| Monroe Cou | 266 | 8.6 (45) | 10.2 (56) | 1,127 | 4.5 (27) | 1.5 (7) | 14.7 (67) | 484 | 15.9 (35) | 12.6 (38) | 818 | 6.6 (31) |
| N.E.W. East | 684 | 8.5 (43) | 7.6 (24) | 2,975 | 4.2 (20) | 2.9 (47) | 13.7 (56) | 1,426 | 17.6 (55) | 16.8 (72) | 2,285 | 3.2 (4) |
| N.E.W. West | 792 | 6.9 (22) | 9.2 (45) | 2,944 | 4.9 (40) | 5.7 (89) | 12.7 (46) | 1,296 | 8.1 (1) | 6.8 (4) | 2,152 | 10.5 (61) |
| Neenah | 339 | 11.8 (66) | 7.7 (25) | 1,249 | 5.0 (43) | 3.6 (62) | 14.7 (67) | 495 | 18.6 (66) | 14.5 (58) | 871 | 12.7 (72) |
| Oconto Cty | 136 | 8.8 (47) | 10.3 (57) | 623 | 4.5 (27) | 1.6 (9) | 15.9 (77) | 287 | 18.8 (68) | 18.1 (76) | 469 | 7.2 (34) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 7
Run Date: 3 / 13 / 09

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | |
|---|-----------------|------------------------|------------------------|-------------------------|-------------------------|---------------------|----------------------|--------------------------------|----------------------------|-----------------------------|--------------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | High | | Short Stature (6) | Under- weight (7) | Obese (7) | | ≥ 2 Yrs Over- weight (7) | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- <95th (Rank) | % ≥95th (Rank) | Number % (Rank) |
| Oneida Cty | 163 | 8.6 (45) | 9.8 (54) | 627 | 5.4 (52) | 4.6 (82) | 9.6 (8) | 267 | 10.9 (6) | 13.1 (43) | 430 5.8 (27) |
| Oneida Tri | 143 | 6.3 (15) | 13.3 (65) | 559 | 5.4 (52) | 2.5 (31) | 22.2 (92) | 231 | 22.1 (83) | 32.0 (86) | 408 3.2 (4) |
| Oshkosh | 338 | 8.3 (40) | 5.9 (11) | 1,320 | 7.2 (88) | 3.0 (50) | 14.5 (65) | 549 | 13.7 (19) | 12.6 (38) | 948 10.1 (57) |
| Outagamie | 677 | 9.3 (51) | 6.5 (16) | 2,523 | 4.0 (17) | 5.0 (84) | 11.8 (34) | 1,163 | 18.5 (64) | 13.2 (44) | 1,655 9.0 (51) |
| Ozaukee | 133 | 6.8 (21) | 13.5 (66) | 487 | 6.0 (65) | 2.9 (47) | 11.1 (23) | 209 | 12.4 (10) | 6.7 (3) | 343 9.9 (56) |
| Pepin | 27 | * * | * * | 150 | 2.7 (7) | 4.7 (83) | 11.3 (25) | 82 | * * | * * | 119 13.4 (73) |
| Pierce Cty | 154 | 7.1 (26) | 10.4 (58) | 711 | 6.8 (82) | 2.0 (17) | 14.3 (64) | 336 | 19.3 (70) | 9.8 (10) | 538 12.3 (69) |
| Polk Cty | 259 | 6.9 (22) | 5.4 (9) | 956 | 3.9 (15) | 4.3 (76) | 9.3 (4) | 400 | 13.3 (17) | 10.0 (12) | 679 9.4 (54) |
| Portage Ct | 278 | 7.6 (32) | 6.8 (19) | 1,187 | 6.1 (69) | 2.7 (38) | 11.7 (32) | 502 | 13.5 (18) | 12.4 (32) | 867 4.3 (15) |
| Price Cty | 79 | * * | * * | 466 | 4.3 (22) | 3.0 (50) | 14.6 (66) | 242 | 21.9 (81) | 15.3 (61) | 382 3.4 (9) |
| Racine Hea | 433 | 10.6 (58) | 6.5 (16) | 1,716 | 3.5 (11) | 2.7 (38) | 13.1 (50) | 765 | 14.2 (20) | 11.8 (24) | 1,308 16.3 (82) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 8
Run Date: 3 / 13 / 09

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | |
|---|-----------------|------------------------|------------------------|-------------------------|-------------------------|---------------------|----------------------|--------------------------------|----------------------------|-----------------------------|--------------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | High | | Short Stature (6) | Under- weight (7) | Obese (7) | | ≥ 2 Yrs Over- weight (7) | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- <95th (Rank) | % ≥95th (Rank) | Number % (Rank) |
| Richland | 99 | * * | * * | 396 | 4.8 (36) | 2.8 (43) | 17.2 (82) | 174 | 19.5 (73) | 20.7 (82) | 278 14.0 (78) |
| Rusk Cty | 85 | * * | * * | 409 | 3.7 (12) | 2.7 (38) | 15.2 (73) | 191 | 22.5 (84) | 19.9 (80) | 307 5.2 (23) |
| Sauk Cty | 337 | 4.5 (5) | 9.2 (45) | 1,371 | 5.0 (43) | 2.3 (23) | 14.2 (63) | 597 | 18.3 (62) | 16.2 (68) | 1,001 2.7 (3) |
| Sawyer Cty | 99 | * * | * * | 451 | 4.9 (40) | 2.4 (28) | 12.2 (37) | 214 | 17.3 (49) | 15.4 (62) | 344 5.2 (23) |
| Seeds Of H | 768 | 7.8 (35) | 8.6 (40) | 3,210 | 3.1 (10) | 4.1 (74) | 12.6 (44) | 1,416 | 18.9 (69) | 15.4 (62) | 2,402 11.1 (64) |
| Shawano Ct | 184 | 6.5 (17) | 6.0 (12) | 812 | 6.4 (74) | 2.2 (19) | 15.1 (72) | 366 | 14.2 (20) | 14.2 (55) | 631 6.3 (30) |
| Sheboygan | 568 | 8.8 (47) | 6.9 (21) | 2,326 | 5.5 (57) | 5.4 (87) | 12.5 (42) | 1,074 | 15.6 (31) | 14.2 (55) | 1,721 3.2 (4) |
| St. Croix | 288 | 6.9 (22) | 8.0 (29) | 1,174 | 6.0 (65) | 2.5 (31) | 11.3 (25) | 521 | 15.7 (33) | 14.4 (57) | 849 4.9 (17) |
| SWCAP | 54 | * * | * * | 217 | 7.8 (89) | 1.8 (14) | 15.2 (73) | 92 | * * | * * | 157 10.2 (59) |
| Taylor Cty | 120 | 10.8 (62) | 11.7 (62) | 443 | 4.1 (19) | 6.5 (91) | 10.4 (14) | 185 | 9.7 (3) | 8.1 (6) | 325 2.5 (2) |
| Trempealeau | 196 | 7.7 (33) | 9.2 (45) | 770 | 4.4 (25) | 2.2 (19) | 17.7 (83) | 336 | 17.6 (55) | 17.6 (75) | 555 9.2 (52) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | | | |
|--|-----------------|------------------------|------------------------|-------------------------|---------------------|-------------------------|----------------------|--------------------------------|----------------------|----------------------|-----------------------------|--------|-------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | | |
| | Low | High | | Short Stature (6) | | Under- weight (7) | Obese (7) | ≥ 2 Yrs Over- weight (7) | | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- (Rank) | % <95th (Rank) | % ≥95th (Rank) | Number | % (Rank) |
| Vernon Cou | 182 | 6.6 (18) | 11.0 (60) | 742 | 5.4 (52) | 2.4 (28) | 12.7 (46) | 327 | 17.7 (58) | 15.9 (66) | | 534 | 5.1 (20) |
| Vilas Cty | 65 | * * | * * | 274 | 6.6 (77) | 5.1 (85) | 10.2 (12) | 114 | 13.2 (15) | 11.4 (21) | | 185 | 3.8 (11) |
| Walworth C | 520 | 5.8 (11) | 8.1 (32) | 2,022 | 5.9 (63) | 2.5 (31) | 14.8 (69) | 828 | 15.2 (26) | 16.2 (68) | | 1,453 | 7.2 (34) |
| Washburn C | 92 | * * | * * | 445 | 4.3 (22) | 7.4 (95) | 9.4 (6) | 210 | 10.5 (5) | 12.4 (32) | | 339 | 8.3 (44) |
| Washington | 401 | 6.2 (13) | 6.2 (13) | 1,494 | 5.8 (62) | 2.3 (23) | 12.2 (37) | 653 | 15.5 (30) | 11.9 (26) | | 1,054 | 8.2 (40) |
| Waukesha C | 651 | 7.5 (29) | 7.8 (28) | 2,400 | 4.8 (36) | 2.3 (23) | 13.7 (56) | 963 | 17.3 (49) | 13.0 (42) | | 1,550 | 6.1 (29) |
| Waupaca Ct | 218 | 10.6 (58) | 8.7 (41) | 830 | 6.5 (75) | 3.9 (71) | 14.9 (70) | 339 | 15.0 (25) | 11.8 (24) | | 591 | 3.4 (9) |
| Waushara | 111 | 4.5 (5) | 12.6 (64) | 488 | 6.6 (77) | 3.1 (52) | 16.4 (81) | 222 | 18.5 (64) | 19.4 (78) | | 357 | 10.4 (60) |
| WC Capitol | 1,021 | 9.6 (54) | 5.2 (8) | 3,807 | 5.0 (43) | 4.2 (75) | 11.0 (20) | 1,587 | 12.7 (13) | 9.0 (8) | | 2,579 | 18.6 (84) |
| WC North | 542 | 12.2 (69) | 3.7 (2) | 2,157 | 4.9 (40) | 4.5 (81) | 9.6 (8) | 952 | 12.5 (11) | 11.9 (26) | | 1,550 | 19.8 (87) |
| WC Teutonia | 537 | 13.4 (71) | 1.7 (1) | 2,298 | 8.0 (92) | 3.7 (64) | 13.4 (53) | 1,066 | 17.0 (47) | 12.9 (40) | | 1,730 | 25.4 (90) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 10
Run Date: 3 / 13 / 09

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | |
|---|------------------|------------------------|------------------------|-------------------------|-------------------------|---------------------|----------------------|--------------------------------|----------------------------|-----------------------------|-----------------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | High | | Short Stature (6) | Under- weight (7) | Obese (7) | | ≥ 2 Yrs Over- weight (7) | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- <95th (Rank) | % ≥95th (Rank) | Number % (Rank) |
| West Allis | 363 | 10.5 (57) | 9.6 (51) | 1,313 | 5.2 (49) | 4.3 (76) | 10.4 (14) | 513 | 17.3 (49) | 10.7 (15) | 892 13.9 (76) |
| Wood Cty | 456 | 7.5 (29) | 7.7 (25) | 1,499 | 5.2 (49) | 3.2 (54) | 11.3 (25) | 594 | 15.8 (34) | 11.1 (20) | 963 7.6 (37) |
| Wisconsin | 31,940 | 8.5 | 7.6 | 126,769 | 5.5 | 3.5 | 13.3 | 55,875 | 16.3 | 13.6 | 91,619 11.0 |
| Nation (Prior Year) | 1,953,242 | 9.1 | 6.4 | 7,526,585 | 6.2 | 4.5 | 13.9 | 3,058,130 | 16.4 | 14.9 | 4,759,588 15.0 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 1
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | | | | Exclusive Breastfeeding (6) | | | | ≤2 Hours TV Viewing/Day (7) | | Smoking in Household | |
|--------------|----------------------|------------------------|--------|--|--------|---|--------------------------------|--|--------|--|-----------------------------------|-----------|-------------------------|-----------|
| Clinic | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months | Number | % Breastfed At Least 3 Months | Number | % Breastfed At Least 6 Months | Number | % | Number | % |
| | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | |
| 16Th St Co | 1,447 | 84.5 (1) | 728 | 43.5 (2) | 1,089 | 26.3 (1) | 497 | 12.7 (21) | 670 | 5.5 (23) | 3,774 | 80.7 (58) | 6,977 | 4.6 (5) |
| Adams Cty | 142 | 64.1 (43) | 56 | * * | 86 | * * | 45 | * * | 54 | * * | 289 | 73.7 (81) | 582 | 20.8 (93) |
| Ashland Cty | 112 | 78.6 (2) | 65 | * * | 91 | * * | 44 | * * | 63 | * * | 280 | 88.6 (11) | 541 | 7.8 (33) |
| Aurora | 1,544 | 38.1 (71) | 369 | 16.5 (43) | 516 | 8.7 (44) | 224 | 10.3 (25) | 332 | 1.8 (43) | 1,545 | 83.2 (42) | 4,110 | 14.7 (78) |
| Barron Cty | 284 | 71.1 (18) | 116 | 28.4 (18) | 160 | 8.8 (42) | 109 | 21.1 (10) | 111 | 8.1 (12) | 522 | 81.4 (55) | 1,137 | 7.8 (33) |
| Bayfield Cty | 55 | * * | 36 | * * | 34 | * * | 22 | * * | 34 | * * | 130 | 87.7 (16) | 243 | 5.8 (14) |
| Beloit | 502 | 48.2 (70) | 215 | 22.8 (35) | 348 | 10.1 (33) | 129 | 10.1 (26) | 204 | 2.5 (39) | 930 | 83.4 (40) | 1,996 | 5.4 (11) |
| Buffalo Cty | 67 | * * | 31 | * * | 41 | * * | 22 | * * | 30 | * * | 126 | 88.9 (10) | 266 | 10.2 (59) |
| Burnett Cty | 88 | * * | 35 | * * | 56 | * * | 32 | * * | 35 | * * | 176 | 98.3 (1) | 384 | 12.2 (73) |
| Calumet Ct | 148 | 62.2 (47) | 67 | * * | 90 | * * | 48 | * * | 63 | * * | 330 | 78.5 (66) | 643 | 9.0 (51) |
| Chippewa C | 335 | 62.1 (48) | 132 | 27.3 (22) | 226 | 11.1 (26) | 79 | * * | 106 | 5.7 (22) | 709 | 88.3 (15) | 1,416 | 8.2 (40) |
| Clark Cty | 180 | 77.2 (6) | 73 | * * | 101 | 6.9 (53) | 54 | * * | 68 | * * | 413 | 86.7 (19) | 803 | 8.1 (37) |
| Columbia C | 221 | 70.1 (21) | 106 | 34.0 (7) | 137 | 16.8 (7) | 75 | * * | 91 | * * | 365 | 79.7 (63) | 853 | 8.9 (49) |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 2
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | Exclusive Breastfeeding (6) | | ≤2 Hours TV Viewing/Day (7) | Smoking in Household |
|--------------|--|--|---|--|--|-----------------------------------|-------------------------------|
| Clinic | Number % Ever Breastfed (Rank) | Number % Breastfed At Least 6 Months (Rank) | Number % Breastfed At Least 12 Months (Rank) | Number % Breastfed At Least 3 Months (Rank) | Number % Breastfed At Least 6 Months (Rank) | Number % (Rank) | Number % (Rank) |
| Crawford Cty | 87 * * | 40 * * | 49 * * | 34 * * | 38 * * | 138 88.4 (14) | 307 3.9 (2) |
| Cudahy | 429 67.1 (32) | 177 25.4 (27) | 274 9.9 (36) | 105 7.6 (35) | 151 3.3 (33) | 696 65.1 (86) | 1,587 18.1 (87) |
| Dane Cty | 0 * * | 0 * * | 3 * * | 0 * * | 0 * * | 0 * * | 6 * * |
| Dodge Cty | 333 76.0 (11) | 136 30.1 (16) | 199 16.6 (8) | 109 22.9 (8) | 131 12.2 (6) | 653 90.5 (5) | 1,340 5.3 (8) |
| Door Cty | 122 67.2 (31) | 51 * * | 71 * * | 36 * * | 49 * * | 239 86.6 (20) | 470 5.1 (7) |
| Douglas Cty | 302 66.2 (39) | 138 30.4 (15) | 205 12.2 (17) | 126 31.0 (1) | 134 20.1 (1) | 570 82.5 (49) | 1,215 8.1 (37) |
| Dunn Cty | 205 73.7 (12) | 102 28.4 (18) | 125 8.8 (42) | 72 * * | 99 * * | 460 85.9 (24) | 920 5.5 (12) |
| Eau Claire | 577 69.8 (22) | 243 25.9 (24) | 345 8.4 (46) | 180 16.7 (14) | 229 7.4 (16) | 1,163 83.1 (45) | 2,394 4.8 (6) |
| Florence C | 25 * * | 14 * * | 14 * * | 9 * * | 12 * * | 67 * * | 127 10.2 (59) |
| Fond Du La | 454 67.4 (30) | 232 29.7 (17) | 329 11.6 (21) | 175 21.7 (9) | 224 6.3 (19) | 974 85.6 (28) | 1,978 7.6 (31) |
| Forest Cty | 47 * * | 22 * * | 33 * * | 14 * * | 22 * * | 116 75.9 (74) | 221 15.4 (84) |
| GLITC-Bad | 22 * * | 10 * * | 18 * * | 7 * * | 10 * * | 63 * * | 118 22.0 (95) |
| GLITC-Bara | 11 * * | 9 * * | 7 * * | 5 * * | 7 * * | 23 * * | 56 * * |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 3
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | Exclusive Breastfeeding (6) | | ≤2 Hours TV Viewing/Day (7) | Smoking in Household |
|---------------|--|--|---|--|--|-----------------------------------|-------------------------------|
| Clinic | Number % Ever Breastfed (Rank) | Number % Breastfed At Least 6 Months (Rank) | Number % Breastfed At Least 12 Months (Rank) | Number % Breastfed At Least 3 Months (Rank) | Number % Breastfed At Least 6 Months (Rank) | Number % (Rank) | Number % (Rank) |
| GLITC-BRF | 18 * * | 19 * * | 20 * * | 8 * * | 17 * * | 62 * * | 112 21.4 (94) |
| GLITC-CEX | 6 * * | 4 * * | 5 * * | 1 * * | 4 * * | 18 * * | 32 * * |
| GLITC-LCO | 35 * * | 10 * * | 25 * * | 2 * * | 9 * * | 72 * * | 173 5.8 (14) |
| GLITC-Mole | 6 * * | 2 * * | 5 * * | 2 * * | 2 * * | 24 * * | 39 * * |
| GLITC-Pete | 50 * * | 32 * * | 47 * * | 20 * * | 29 * * | 149 79.9 (62) | 283 19.8 (91) |
| GLITC-Pota | 16 * * | 5 * * | 11 * * | 1 * * | 3 * * | 39 * * | 81 * * |
| GLITC-Red | 27 * * | 11 * * | 15 * * | 9 * * | 10 * * | 65 * * | 111 16.2 (86) |
| GLITC-St.C | 29 * * | 10 * * | 17 * * | 7 * * | 9 * * | 65 * * | 156 4.5 (4) |
| GLITC-Stoc | 25 * * | 13 * * | 13 * * | 5 * * | 13 * * | 75 * * | 137 8.8 (47) |
| Grant Cty | 286 60.8 (51) | 140 26.4 (23) | 161 12.4 (16) | 79 * * | 114 4.4 (27) | 617 70.5 (82) | 1,240 8.2 (40) |
| Green Cty | 148 58.1 (59) | 82 * * | 105 10.5 (29) | 62 * * | 80 * * | 319 87.5 (17) | 648 12.0 (72) |
| Green Lake | 69 * * | 38 * * | 51 * * | 24 * * | 31 * * | 146 75.3 (76) | 310 10.0 (56) |
| Greenfield | 273 71.1 (18) | 97 * * | 150 10.0 (35) | 72 * * | 89 * * | 360 75.8 (75) | 882 15.0 (81) |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 4
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | | | | Exclusive Breastfeeding (6) | | | | ≤2 Hours TV Viewing/Day (7) | | Smoking in Household | |
|------------|----------------------|------------------------|--------|--|--------|---|--------------------------------|--|--------|--|-----------------------------------|-----------|-------------------------|-----------|
| Clinic | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months | Number | % Breastfed At Least 3 Months | Number | % Breastfed At Least 6 Months | Number | % | Number | % |
| | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | |
| Hlth & Nut | 1,023 | 61.3 (50) | 373 | 23.9 (33) | 525 | 10.3 (31) | 300 | 12.3 (22) | 359 | 4.7 (25) | 1,667 | 80.0 (61) | 3,627 | 10.1 (58) |
| Iowa Cty | 110 | 65.5 (41) | 50 | * * | 58 | * * | 39 | * * | 46 | * * | 140 | 90.0 (6) | 354 | 10.2 (59) |
| Iron Cty | 27 | * * | 8 | * * | 12 | * * | 11 | * * | 8 | * * | 46 | * * | 96 | * * |
| Jackson Ct | 119 | 66.4 (36) | 50 | * * | 79 | * * | 41 | * * | 46 | * * | 298 | 82.2 (51) | 569 | 7.9 (35) |
| Janesville | 571 | 49.7 (69) | 223 | 18.8 (41) | 304 | 11.5 (23) | 184 | 9.2 (29) | 205 | 3.9 (29) | 910 | 84.8 (31) | 2,039 | 11.7 (68) |
| Jefferson | 406 | 68.2 (25) | 203 | 25.1 (28) | 258 | 11.6 (21) | 144 | 6.9 (37) | 178 | 3.4 (32) | 817 | 86.9 (18) | 1,693 | 8.2 (40) |
| Juneau | 158 | 72.2 (14) | 85 | * * | 112 | 8.0 (47) | 77 | * * | 83 | * * | 348 | 76.1 (73) | 720 | 19.3 (90) |
| Kenosha | 1,100 | 62.3 (46) | 429 | 21.2 (38) | 651 | 8.6 (45) | 316 | 7.6 (35) | 394 | 2.8 (35) | 1,794 | 80.9 (57) | 4,073 | 9.4 (52) |
| Kewaunee C | 77 | * * | 32 | * * | 39 | * * | 29 | * * | 29 | * * | 158 | 67.7 (84) | 323 | 7.1 (28) |
| La Crosse | 595 | 67.6 (29) | 259 | 24.7 (30) | 341 | 16.4 (10) | 170 | 20.6 (11) | 208 | 6.7 (18) | 1,141 | 86.6 (20) | 2,415 | 3.8 (1) |
| Langlade | 113 | 57.5 (60) | 54 | * * | 68 | * * | 35 | * * | 48 | * * | 240 | 83.3 (41) | 488 | 15.0 (81) |
| Lincoln | 110 | 56.4 (62) | 52 | * * | 76 | * * | 35 | * * | 46 | * * | 228 | 89.0 (9) | 474 | 13.7 (77) |
| Madison WI | 1,885 | 76.7 (8) | 890 | 45.1 (1) | 1,167 | 22.9 (2) | 737 | 19.4 (13) | 804 | 10.9 (8) | 3,306 | 84.7 (32) | 7,194 | 5.9 (17) |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 5
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | | | | Exclusive Breastfeeding (6) | | | | ≤2 Hours TV Viewing/Day (7) | | Smoking in Household | |
|-------------|----------------------|------------------------|--------|--|--------|---|--------------------------------|--|--------|--|-----------------------------------|-----------|-------------------------|-----------|
| Clinic | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months | Number | % Breastfed At Least 3 Months | Number | % Breastfed At Least 6 Months | Number | % | Number | % |
| | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | |
| Manitowoc | 402 | 60.0 (52) | 154 | 24.0 (32) | 257 | 11.7 (20) | 118 | 23.7 (7) | 147 | 8.2 (11) | 786 | 76.8 (70) | 1,649 | 8.6 (45) |
| Marathon | 625 | 59.0 (55) | 324 | 21.9 (36) | 382 | 9.4 (40) | 207 | 15.9 (15) | 287 | 5.9 (20) | 1,274 | 82.3 (50) | 2,673 | 7.2 (29) |
| Marinette | 222 | 50.0 (68) | 110 | 21.8 (37) | 160 | 3.1 (57) | 72 | * * | 104 | 11.5 (7) | 525 | 86.3 (22) | 1,046 | 11.4 (67) |
| Marquette | 59 | * * | 22 | * * | 41 | * * | 26 | * * | 16 | * * | 123 | 74.8 (79) | 247 | 15.0 (81) |
| Menominee | 146 | 66.4 (36) | 57 | * * | 94 | * * | 32 | * * | 54 | * * | 289 | 89.3 (8) | 555 | 7.6 (31) |
| MilwaukeeJ | 591 | 77.5 (4) | 248 | 37.1 (3) | 360 | 19.2 (3) | 186 | 9.7 (27) | 235 | 2.6 (38) | 1,343 | 81.6 (54) | 2,580 | 8.2 (40) |
| MilwaukeeK | 506 | 51.8 (66) | 223 | 13.0 (46) | 411 | 4.6 (56) | 124 | 8.9 (30) | 217 | 2.3 (40) | 1,173 | 66.4 (85) | 2,324 | 20.1 (92) |
| MilwaukeeN | 903 | 58.5 (56) | 383 | 11.5 (47) | 596 | 7.0 (51) | 239 | 5.9 (39) | 375 | 1.9 (42) | 1,643 | 76.6 (72) | 3,644 | 10.7 (63) |
| MilwaukMLK | 632 | 57.4 (61) | 350 | 10.6 (48) | 547 | 6.9 (53) | 203 | 7.9 (34) | 332 | 2.7 (36) | 1,378 | 74.2 (80) | 2,880 | 18.4 (89) |
| Monroe Cou | 272 | 69.5 (23) | 122 | 32.8 (11) | 182 | 13.7 (15) | 90 | * * | 109 | 12.8 (4) | 521 | 85.6 (28) | 1,142 | 10.0 (56) |
| N.E.W. East | 680 | 62.4 (45) | 322 | 24.5 (31) | 506 | 11.9 (19) | 237 | 9.7 (27) | 297 | 3.7 (31) | 1,550 | 77.8 (69) | 3,045 | 5.5 (12) |
| N.E.W. West | 796 | 55.5 (63) | 352 | 23.6 (34) | 493 | 9.5 (39) | 230 | 13.9 (19) | 305 | 3.9 (29) | 1,369 | 79.3 (64) | 2,969 | 11.0 (66) |
| Neenah | 338 | 67.8 (27) | 153 | 24.8 (29) | 184 | 9.2 (41) | 113 | 15.9 (15) | 147 | 7.5 (15) | 537 | 83.2 (42) | 1,223 | 6.7 (22) |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 6
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | | | | Exclusive Breastfeeding (6) | | ≤2 Hours TV Viewing/Day (7) | | Smoking in Household | | | |
|------------|----------------------|------------------------|--------|--|--------|---|--------------------------------|--|-----------------------------------|--|-------------------------|-----------|-------|-----------|
| Clinic | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months | Number | % Breastfed At Least 3 Months | Number | % Breastfed At Least 6 Months | Number | % | | |
| | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | | |
| Oconto Cty | 136 | 59.6 (53) | 75 | * * | 91 | * * | 42 | * * | 68 | * * | 308 | 84.7 (32) | 631 | 12.8 (75) |
| Oneida Cty | 163 | 58.3 (58) | 69 | * * | 80 | * * | 59 | * * | 67 | * * | 283 | 91.9 (3) | 631 | 8.7 (46) |
| Oneida Tri | 143 | 61.5 (49) | 80 | * * | 108 | 7.4 (50) | 57 | * * | 78 | * * | 254 | 81.1 (56) | 576 | 18.1 (87) |
| Oshkosh | 340 | 58.5 (56) | 136 | 20.6 (39) | 201 | 7.5 (49) | 101 | 13.9 (19) | 134 | 3.0 (34) | 578 | 84.1 (36) | 1,282 | 6.9 (26) |
| Outagamie | 678 | 66.5 (35) | 250 | 28.0 (20) | 363 | 9.9 (36) | 221 | 19.5 (12) | 246 | 7.7 (13) | 1,251 | 82.1 (52) | 2,575 | 5.9 (17) |
| Ozaukee | 134 | 76.9 (7) | 58 | * * | 72 | * * | 62 | * * | 56 | * * | 225 | 83.6 (37) | 489 | 5.3 (8) |
| Pepin | 27 | * * | 17 | * * | 20 | * * | 12 | * * | 17 | * * | 84 | * * | 149 | 13.4 (76) |
| Pierce Cty | 155 | 73.5 (13) | 83 | * * | 103 | 10.7 (28) | 70 | * * | 78 | * * | 360 | 90.8 (4) | 716 | 10.8 (64) |
| Polk Cty | 259 | 76.1 (10) | 121 | 36.4 (6) | 149 | 10.1 (33) | 95 | * * | 111 | 12.6 (5) | 434 | 82.0 (53) | 977 | 8.1 (37) |
| Portage Ct | 275 | 72.0 (16) | 117 | 31.6 (13) | 159 | 13.8 (14) | 99 | * * | 108 | 10.2 (9) | 549 | 88.5 (13) | 1,198 | 6.7 (22) |
| Price Cty | 79 | * * | 52 | * * | 66 | * * | 27 | * * | 41 | * * | 259 | 82.6 (48) | 475 | 9.7 (53) |
| Racine Hea | 433 | 54.3 (64) | 186 | 18.8 (41) | 261 | 10.3 (31) | 113 | 10.6 (24) | 182 | 2.7 (36) | 843 | 78.4 (67) | 1,777 | 8.0 (36) |
| Richland | 99 | * * | 40 | * * | 55 | * * | 31 | * * | 36 | * * | 182 | 83.0 (46) | 396 | 6.1 (19) |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 7
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | Exclusive Breastfeeding (6) | | ≤2 Hours TV Viewing/Day (7) | | Smoking in Household | |
|-------------|--|--|---|--|--|-----------------------------------|--|-------------------------|--|
| Clinic | Number % Ever Breastfed (Rank) | Number % Breastfed At Least 6 Months (Rank) | Number % Breastfed At Least 12 Months (Rank) | Number % Breastfed At Least 3 Months (Rank) | Number % Breastfed At Least 6 Months (Rank) | Number % | | Number % | |
| Rusk Cty | 85 * * | 42 * * | 72 * * | 23 * * | 37 * * | 204 84.3 (35) | | 414 11.8 (70) | |
| Sauk Cty | 335 77.9 (3) | 173 37.0 (4) | 196 17.3 (5) | 171 25.1 (5) | 168 7.7 (13) | 646 79.3 (64) | | 1,389 10.9 (65) | |
| Sawyer Cty | 99 * * | 51 * * | 63 * * | 39 * * | 49 * * | 227 80.6 (59) | | 460 11.7 (68) | |
| Seeds Of H | 768 68.1 (26) | 356 27.8 (21) | 612 12.1 (18) | 246 8.9 (30) | 341 5.0 (24) | 1,562 83.2 (42) | | 3,167 6.8 (25) | |
| Shawano Ct | 184 66.8 (34) | 87 * * | 143 10.5 (29) | 54 * * | 82 * * | 405 85.7 (27) | | 832 9.9 (55) | |
| Sheboygan | 567 64.6 (42) | 239 25.9 (24) | 403 11.4 (24) | 208 14.4 (18) | 228 8.3 (10) | 1,150 83.0 (46) | | 2,376 8.9 (49) | |
| St. Croix | 292 76.4 (9) | 148 32.4 (12) | 165 18.8 (4) | 103 30.1 (3) | 145 14.5 (2) | 546 80.6 (59) | | 1,179 5.3 (8) | |
| SWCAP | 54 * * | 23 * * | 33 * * | 17 * * | 23 * * | 99 * * | | 218 6.9 (26) | |
| Taylor Cty | 120 68.3 (24) | 47 * * | 65 * * | 32 * * | 42 * * | 202 78.2 (68) | | 450 11.8 (70) | |
| Trempealeau | 195 67.7 (28) | 99 * * | 108 15.7 (11) | 80 * * | 92 * * | 371 95.7 (2) | | 795 4.4 (3) | |
| Vernon Cou | 182 72.0 (16) | 79 * * | 116 17.2 (6) | 57 * * | 71 * * | 360 85.8 (26) | | 751 7.3 (30) | |
| Vilas Cty | 65 * * | 31 * * | 43 * * | 29 * * | 29 * * | 127 83.5 (38) | | 272 8.8 (47) | |
| Walworth C | 521 77.4 (5) | 289 37.0 (4) | 283 16.6 (8) | 229 24.5 (6) | 273 5.9 (20) | 906 84.7 (32) | | 2,050 5.8 (14) | |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 8
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | Exclusive Breastfeeding (6) | | | ≤2 Hours TV Viewing/Day (7) | | Smoking in Household | |
|--------|--|--|--|--------------------------------|--|--|-----------------------------------|--|-------------------------|--|
| Clinic | Number | | | | | | | | | |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | | | |
|--|-----------------|------------------------|------------------------|-------------------------|---------------------|-------------------------|----------------------|--------------------------------|----------------------|----------------------|-----------------------------|--------|-------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | | |
| | Low | | High | Short Stature (6) | | Under- weight (7) | Obese (7) | ≥ 2 Yrs Over- weight (7) | | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- (Rank) | % <95th (Rank) | % ≥95th (Rank) | Number | % (Rank) |
| 16Th St Co | 4,364 | 6.4 (17) | 8.2 (40) | 20,217 | 7.1 (94) | 1.5 (5) | 19.8 (93) | 9,798 | 20.8 (87) | 22.4 (90) | | 15,497 | 9.9 (63) |
| Adams Cty | 352 | 10.8 (82) | 9.4 (61) | 1,500 | 5.9 (72) | 3.1 (45) | 14.7 (78) | 683 | 17.6 (59) | 13.8 (57) | | 1,091 | 8.0 (42) |
| Ashland Cty | 351 | 5.4 (6) | 12.5 (83) | 1,606 | 5.9 (72) | 4.0 (70) | 9.7 (9) | 766 | 11.4 (6) | 8.5 (7) | | 1,244 | 5.2 (23) |
| Aurora | 4,433 | 9.2 (71) | 4.2 (5) | 11,755 | 4.5 (26) | 7.1(100) | 9.0 (4) | 4,266 | 13.9 (16) | 10.5 (18) | | 6,510 | 28.8 (99) |
| Barron Cty | 835 | 8.4 (55) | 9.5 (63) | 3,236 | 5.3 (52) | 2.4 (21) | 12.6 (52) | 1,361 | 17.4 (58) | 14.1 (63) | | 2,281 | 8.3 (47) |
| Bayfield Cty | 166 | 7.2 (27) | 12.7 (84) | 724 | 4.3 (19) | 4.6 (82) | 9.9 (14) | 353 | 21.2 (89) | 9.9 (14) | | 553 | 5.6 (28) |
| Beloit | 1,567 | 9.1 (70) | 6.8 (15) | 5,867 | 5.5 (61) | 3.2 (52) | 14.8 (80) | 2,457 | 16.4 (44) | 15.1 (73) | | 4,101 | 9.3 (59) |
| Buffalo Cty | 197 | 8.6 (62) | 6.6 (13) | 794 | 2.9 (3) | 2.6 (25) | 9.8 (12) | 353 | 18.4 (71) | 13.3 (54) | | 564 | 12.6 (80) |
| Burnett Cty | 276 | 8.3 (52) | 7.6 (27) | 1,145 | 5.3 (52) | 4.5 (81) | 8.4 (1) | 521 | 14.4 (19) | 7.5 (3) | | 838 | 1.9 (2) |
| Calumet Ct | 448 | 9.8 (77) | 7.8 (32) | 1,840 | 6.6 (90) | 4.0 (70) | 12.1 (45) | 850 | 17.1 (53) | 14.7 (70) | | 1,385 | 4.6 (20) |
| Chippewa C | 1,005 | 7.5 (34) | 7.4 (24) | 4,092 | 6.2 (82) | 3.1 (45) | 11.0 (25) | 1,838 | 15.6 (32) | 11.5 (31) | | 3,022 | 8.5 (51) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | | |
|--|-----------------|------------------------|------------------------|-------------------------|---------------------|---------------------|-------------------------|--------------|--------------------------------|----------------------|-----------------------------|-------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | | High | Short Stature (6) | | | Under- weight (7) | Obese (7) | ≥ 2 Yrs Over- weight (7) | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- <95th (Rank) | % ≥95th (Rank) | Number | % (Rank) |
| Clark Cty | 532 | 11.1 (84) | 7.7 (29) | 2,279 | 5.4 (57) | 3.0 (40) | 14.1 (70) | 1,026 | 16.9 (49) | 16.2 (80) | 1,670 | 13.5 (84) |
| Columbia C | 640 | 9.2 (71) | 9.7 (70) | 2,474 | 6.0 (76) | 3.7 (62) | 11.6 (35) | 1,032 | 16.0 (39) | 14.0 (62) | 1,723 | 6.7 (35) |
| Crawford Cty | 244 | 7.8 (40) | 8.2 (40) | 914 | 6.6 (90) | 3.1 (45) | 10.7 (21) | 389 | 15.2 (28) | 11.3 (27) | 630 | 9.8 (62) |
| Cudahy | 1,227 | 9.3 (73) | 9.4 (61) | 4,241 | 5.4 (57) | 3.6 (60) | 11.1 (27) | 1,638 | 15.2 (28) | 12.1 (38) | 2,966 | 8.1 (44) |
| Dane Cty | 941 | 7.5 (34) | 9.5 (63) | 3,708 | 5.0 (42) | 3.1 (45) | 10.8 (24) | 1,590 | 15.3 (30) | 12.2 (39) | 2,703 | 8.6 (54) |
| Dodge Cty | 949 | 8.9 (68) | 7.0 (21) | 3,700 | 4.6 (28) | 4.6 (82) | 9.2 (5) | 1,633 | 11.3 (5) | 8.1 (4) | 2,636 | 6.2 (31) |
| Door Cty | 328 | 5.2 (4) | 8.5 (48) | 1,357 | 3.5 (9) | 2.9 (34) | 9.2 (5) | 644 | 18.9 (75) | 13.2 (50) | 998 | 4.1 (18) |
| Douglas Cty | 823 | 5.1 (2) | 11.5 (81) | 3,500 | 4.1 (15) | 3.9 (68) | 12.5 (50) | 1,589 | 19.1 (78) | 14.5 (67) | 2,378 | 13.9 (85) |
| Dunn Cty | 631 | 6.2 (14) | 10.1 (76) | 2,804 | 6.0 (76) | 4.4 (79) | 11.7 (39) | 1,286 | 17.0 (51) | 13.0 (47) | 2,035 | 3.4 (7) |
| Eau Claire | 1,683 | 6.2 (14) | 8.6 (51) | 6,912 | 5.7 (68) | 2.2 (18) | 10.5 (18) | 3,068 | 15.8 (37) | 10.9 (22) | 5,085 | 11.6 (76) |
| Florence C | 70 | * * | * * | 351 | 7.4 (96) | 1.4 (4) | 10.5 (18) | 180 | 15.0 (25) | 8.3 (6) | 248 | 11.3 (75) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | | |
|--|-----------------|------------------------|------------------------|-------------------------|---------------------|-------------------------|----------------------|--------------------------------|----------------------------|----------------------|-----------------------------|-------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | | High | Short Stature (6) | | Under- weight (7) | Obese (7) | ≥ 2 Yrs Over- weight (7) | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- <95th (Rank) | % ≥95th (Rank) | Number | % (Rank) |
| Fond Du La | 1,394 | 9.0 (69) | 6.5 (12) | 5,662 | 6.4 (87) | 3.4 (56) | 9.2 (5) | 2,563 | 14.6 (21) | 8.9 (8) | 4,192 | 8.9 (56) |
| Forest Cty | 162 | 8.6 (62) | 6.8 (15) | 712 | 6.2 (82) | 3.8 (65) | 15.0 (82) | 351 | 13.7 (14) | 16.0 (79) | 542 | 7.0 (36) |
| GLITC-Bad | 70 | * * | * * | 354 | 3.1 (6) | 1.7 (9) | 20.9 (94) | 186 | 26.3 (98) | 24.7 (94) | 278 | 7.6 (40) |
| GLITC-Bara | 35 | * * | * * | 140 | 8.6(100) | 5.0 (89) | 35.0(101) | 57 | * * | * * | 97 | * * |
| GLITC-BRF | 77 | * * | * * | 365 | 4.4 (22) | 1.6 (6) | 26.0 (99) | 171 | 23.4 (95) | 28.7 (97) | 288 | 16.7 (90) |
| GLITC-CEX | 23 | * * | * * | 91 | * * | * * | * * | 44 | * * | * * | 67 | * * |
| GLITC-LCO | 127 | 3.9 (1) | 15.0 (87) | 529 | 4.3 (19) | 3.0 (40) | 16.3 (86) | 221 | 22.6 (92) | 20.4 (88) | 380 | 14.5 (88) |
| GLITC-Mole | 20 | * * | * * | 102 | 2.9 (3) | 2.0 (14) | 19.6 (92) | 49 | * * | * * | 84 | * * |
| GLITC-Pete | 178 | 7.9 (44) | 13.5 (85) | 770 | 3.8 (13) | 1.9 (12) | 23.4 (97) | 346 | 23.4 (95) | 23.4 (91) | 586 | 3.6 (10) |
| GLITC-Pota | 51 | * * | * * | 236 | 4.7 (31) | 2.5 (23) | 22.9 (96) | 108 | 20.4 (84) | 27.8 (96) | 144 | 16.7 (90) |
| GLITC-Red | 77 | * * | * * | 330 | 4.8 (34) | 7.9(101) | 19.1 (90) | 159 | 17.6 (59) | 23.9 (92) | 245 | 1.2 (1) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | | |
|--|-----------------|------------------------|------------------------|-------------------------|---------------------|-------------------------|----------------------|--------------------------------|----------------------------|----------------------|-----------------------------|-------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | | High | Short Stature (6) | | Under- weight (7) | Obese (7) | ≥ 2 Yrs Over- weight (7) | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- <95th (Rank) | % ≥95th (Rank) | Number | % (Rank) |
| GLITC-St.C | 30 | * * | * * | 153 | 4.6 (28) | 1.3 (2) | 29.4(100) | 79 | * * | * * | 122 | 8.2 (46) |
| GLITC-St.C | 60 | * * | * * | 287 | 2.1 (1) | 3.5 (58) | 24.7 (98) | 127 | 20.5 (85) | 29.9 (98) | 231 | 10.8 (70) |
| GLITC-Stoc | 84 | * * | * * | 407 | 2.9 (3) | 1.2 (1) | 17.0 (88) | 199 | 15.6 (32) | 12.6 (43) | 309 | 9.7 (61) |
| Grant Cty | 801 | 6.9 (20) | 9.7 (70) | 3,475 | 4.2 (17) | 4.2 (74) | 9.8 (12) | 1,590 | 14.8 (22) | 11.4 (28) | 2,596 | 4.7 (21) |
| Green Cty | 472 | 8.5 (59) | 9.7 (70) | 1,911 | 7.8 (98) | 2.8 (29) | 17.4 (89) | 807 | 20.6 (86) | 24.3 (93) | 1,353 | 8.1 (44) |
| Green Lake | 237 | 7.2 (27) | 10.1 (76) | 903 | 3.3 (8) | 4.0 (70) | 13.1 (59) | 382 | 13.4 (12) | 12.8 (45) | 637 | 10.4 (66) |
| Greenfield | 727 | 8.4 (55) | 9.1 (58) | 2,345 | 6.3 (86) | 3.8 (65) | 10.1 (15) | 923 | 15.7 (35) | 10.2 (16) | 1,569 | 12.9 (82) |
| Hlth & Nut | 2,848 | 9.6 (75) | 7.3 (22) | 10,198 | 6.2 (82) | 3.0 (40) | 13.7 (67) | 4,235 | 17.6 (59) | 14.4 (66) | 7,113 | 7.6 (40) |
| Iowa Cty | 303 | 8.6 (62) | 7.6 (27) | 1,016 | 5.3 (52) | 2.8 (29) | 13.1 (59) | 384 | 20.3 (83) | 13.8 (57) | 658 | 8.4 (49) |
| Iron Cty | 62 | * * | * * | 277 | 6.9 (93) | 5.8 (95) | 15.9 (84) | 137 | 21.2 (89) | 13.1 (48) | 210 | 10.5 (68) |
| Jackson Ct | 379 | 5.8 (8) | 11.3 (80) | 1,686 | 5.1 (44) | 3.1 (45) | 11.7 (39) | 781 | 21.6 (91) | 11.4 (28) | 1,155 | 8.0 (42) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | | | |
|--|-----------------|------------------------|------------------------|-------------------------|---------------------|---------------------|-------------------------|--------------|--------------------------------|----------------------|-----------------------------|--------|-------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | | |
| | Low | | High | Short Stature (6) | | | Under- weight (7) | Obese (7) | ≥ 2 Yrs Over- weight (7) | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- (Rank) | % <95th (Rank) | % ≥95th (Rank) | Number | % (Rank) |
| Janesville | 1,631 | 8.4 (55) | 9.3 (60) | 5,762 | 4.8 (34) | 4.3 (76) | 11.7 (39) | 2,280 | 16.1 (41) | 11.8 (36) | | 3,862 | 9.0 (57) |
| Jefferson | 1,217 | 5.9 (9) | 8.4 (44) | 5,081 | 3.5 (9) | 3.3 (55) | 11.8 (43) | 2,211 | 17.0 (51) | 15.7 (78) | | 3,637 | 5.4 (25) |
| Juneau | 563 | 6.0 (10) | 9.2 (59) | 2,295 | 5.6 (64) | 2.1 (15) | 14.6 (74) | 1,016 | 17.9 (66) | 13.2 (50) | | 1,640 | 7.0 (36) |
| Kenosha | 3,212 | 9.7 (76) | 6.1 (11) | 11,689 | 7.6 (97) | 2.4 (21) | 14.7 (78) | 4,847 | 16.2 (42) | 13.9 (60) | | 8,073 | 16.9 (92) |
| Kewaunee C | 215 | 7.9 (44) | 8.4 (44) | 933 | 4.9 (38) | 1.3 (2) | 15.9 (84) | 427 | 18.7 (73) | 14.5 (67) | | 685 | 12.7 (81) |
| La Crosse | 1,741 | 7.3 (30) | 9.6 (66) | 6,786 | 6.4 (87) | 4.0 (70) | 11.3 (31) | 3,061 | 19.0 (77) | 12.4 (40) | | 5,026 | 6.6 (34) |
| Langlade | 347 | 7.5 (34) | 8.9 (55) | 1,386 | 5.2 (49) | 2.1 (15) | 12.6 (52) | 616 | 18.7 (73) | 13.8 (57) | | 1,036 | 11.0 (72) |
| Lincoln | 351 | 7.1 (22) | 8.8 (54) | 1,314 | 6.0 (76) | 1.6 (6) | 14.3 (71) | 598 | 17.1 (53) | 15.2 (74) | | 951 | 2.9 (4) |
| Madison WI | 4,548 | 8.2 (49) | 8.3 (42) | 16,416 | 5.0 (42) | 2.9 (34) | 13.0 (57) | 6,962 | 12.9 (8) | 10.9 (22) | | 11,690 | 11.1 (73) |
| Manitowoc | 1,113 | 8.5 (59) | 7.3 (22) | 4,645 | 5.2 (49) | 2.7 (27) | 13.0 (57) | 2,082 | 19.1 (78) | 14.8 (71) | | 3,299 | 10.1 (65) |
| Marathon | 1,871 | 7.1 (22) | 5.8 (9) | 7,476 | 8.6(100) | 1.8 (10) | 13.2 (62) | 3,232 | 18.4 (71) | 13.2 (50) | | 5,522 | 3.5 (8) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | | | |
|--|-----------------|------------------------|------------------------|-------------------------|---------------------|-------------------------|----------------------|--------------------------------|----------------------|----------------------|-----------------------------|--------|-------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | | |
| | Low | High | | Short Stature (6) | | Under- weight (7) | Obese (7) | ≥ 2 Yrs Over- weight (7) | | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- (Rank) | % <95th (Rank) | % ≥95th (Rank) | Number | % (Rank) |
| Marinette | 688 | 6.8 (18) | 9.6 (66) | 3,094 | 4.7 (31) | 4.8 (86) | 9.7 (9) | 1,414 | 9.0 (2) | 7.1 (2) | | 2,327 | 3.7 (12) |
| Marquette | 164 | 6.1 (12) | 7.9 (35) | 673 | 4.6 (28) | 5.3 (90) | 13.4 (63) | 299 | 14.4 (19) | 12.0 (37) | | 475 | 9.5 (60) |
| Menominee | 421 | 6.9 (20) | 12.1 (82) | 1,640 | 2.1 (1) | 1.6 (6) | 21.0 (95) | 771 | 23.2 (93) | 20.5 (89) | | 1,228 | 3.9 (14) |
| MilwaukeeJ | 1,580 | 7.3 (30) | 7.5 (25) | 7,065 | 5.5 (61) | 3.7 (62) | 13.6 (65) | 3,415 | 16.9 (49) | 14.2 (65) | | 5,479 | 14.0 (86) |
| MilwaukeeK | 1,584 | 14.0 (89) | 4.0 (3) | 6,913 | 5.6 (64) | 5.3 (90) | 10.7 (21) | 3,295 | 15.1 (26) | 11.2 (25) | | 5,312 | 20.5 (95) |
| MilwaukeeN | 2,880 | 11.6 (85) | 4.8 (6) | 10,521 | 7.1 (94) | 5.4 (92) | 11.6 (35) | 4,388 | 16.8 (48) | 12.4 (40) | | 7,563 | 17.4 (93) |
| MilwaukMLK | 2,160 | 13.0 (87) | 3.9 (2) | 8,882 | 5.4 (57) | 6.3 (97) | 10.5 (18) | 3,865 | 11.6 (7) | 8.1 (4) | | 6,600 | 23.4 (96) |
| Monroe Cou | 797 | 7.7 (38) | 8.4 (44) | 3,116 | 5.5 (61) | 2.2 (18) | 14.8 (80) | 1,307 | 15.1 (26) | 12.4 (40) | | 2,190 | 5.5 (27) |
| N.E.W. East | 2,086 | 8.2 (49) | 7.8 (32) | 8,736 | 4.4 (22) | 3.1 (45) | 13.1 (59) | 4,112 | 18.1 (67) | 17.3 (83) | | 6,629 | 3.6 (10) |
| N.E.W. West | 2,246 | 7.5 (34) | 8.3 (42) | 8,408 | 4.9 (38) | 6.3 (97) | 11.2 (28) | 3,792 | 8.5 (1) | 6.4 (1) | | 6,154 | 8.5 (51) |
| Neenah | 945 | 9.8 (77) | 6.9 (20) | 3,534 | 4.9 (38) | 2.9 (34) | 13.8 (68) | 1,463 | 17.1 (53) | 15.5 (77) | | 2,453 | 14.0 (86) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 7
Run Date: 3 / 13 / 09

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | | | |
|--|-----------------|------------------------|------------------------|-------------------------|---------------------|---------------------|-------------------------|--------------|--------------------------------|----------------------|-----------------------------|--------|-------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | | |
| | Low | | High | Short Stature (6) | | | Under- weight (7) | Obese (7) | ≥ 2 Yrs Over- weight (7) | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- (Rank) | % <95th (Rank) | % ≥95th (Rank) | Number | % (Rank) |
| Oconto Cty | 441 | 7.9 (44) | 7.9 (35) | 1,775 | 5.3 (52) | 3.4 (56) | 16.9 (87) | 778 | 17.7 (63) | 19.3 (87) | | 1,283 | 7.5 (39) |
| Oneida Cty | 461 | 8.5 (59) | 8.0 (38) | 1,772 | 4.7 (31) | 4.7 (85) | 8.5 (2) | 729 | 13.7 (14) | 9.2 (11) | | 1,218 | 6.5 (33) |
| Oneida Tri | 433 | 5.3 (5) | 16.9 (89) | 1,608 | 4.2 (17) | 1.9 (12) | 19.2 (91) | 696 | 19.8 (82) | 24.7 (94) | | 1,143 | 2.9 (4) |
| Oshkosh | 1,009 | 9.9 (79) | 7.5 (25) | 3,968 | 6.4 (87) | 3.2 (52) | 14.6 (74) | 1,637 | 15.9 (38) | 11.5 (31) | | 2,837 | 11.8 (78) |
| Outagamie | 1,943 | 9.5 (74) | 6.8 (15) | 7,437 | 5.1 (44) | 4.9 (88) | 11.6 (35) | 3,423 | 18.2 (68) | 14.1 (63) | | 4,984 | 9.1 (58) |
| Ozaukee | 374 | 6.1 (12) | 10.2 (78) | 1,400 | 6.1 (80) | 3.0 (40) | 12.1 (45) | 586 | 13.1 (10) | 9.0 (9) | | 981 | 8.3 (47) |
| Pepin | 81 | * * | * * | 374 | 4.3 (19) | 5.6 (94) | 11.2 (28) | 182 | 20.9 (88) | 15.4 (76) | | 282 | 12.4 (79) |
| Pierce Cty | 476 | 5.7 (7) | 9.5 (63) | 2,097 | 6.2 (82) | 2.1 (15) | 12.5 (50) | 977 | 18.9 (75) | 10.4 (17) | | 1,567 | 10.9 (71) |
| Polk Cty | 721 | 7.8 (40) | 6.8 (15) | 2,606 | 4.1 (15) | 4.3 (76) | 8.9 (3) | 1,076 | 14.2 (18) | 9.1 (10) | | 1,835 | 8.8 (55) |
| Portage Ct | 863 | 7.3 (30) | 9.6 (66) | 3,459 | 6.0 (76) | 2.9 (34) | 11.2 (28) | 1,452 | 14.9 (24) | 11.6 (33) | | 2,429 | 4.5 (19) |
| Price Cty | 256 | 6.3 (16) | 9.0 (57) | 1,374 | 3.6 (12) | 3.1 (45) | 12.3 (47) | 677 | 17.6 (59) | 13.1 (48) | | 1,081 | 3.5 (8) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | | |
|--|-----------------|------------------------|------------------------|-------------------------|---------------------|---------------------|-------------------------|--------------|--------------------------------|----------------------|-----------------------------|-------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | | High | Short Stature (6) | | | Under- weight (7) | Obese (7) | ≥ 2 Yrs Over- weight (7) | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- <95th (Rank) | % ≥95th (Rank) | Number | % (Rank) |
| Racine Hea | 1,251 | 10.1 (80) | 5.7 (8) | 4,848 | 3.8 (13) | 2.9 (34) | 12.4 (48) | 2,159 | 14.8 (22) | 11.2 (25) | 3,576 | 16.3 (89) |
| Richland | 255 | 5.1 (2) | 15.3 (88) | 1,024 | 4.9 (38) | 2.6 (25) | 14.4 (72) | 453 | 23.2 (93) | 17.4 (84) | 716 | 11.2 (74) |
| Rusk Cty | 288 | 8.0 (47) | 11.1 (79) | 1,280 | 4.4 (22) | 1.8 (10) | 14.6 (74) | 610 | 18.2 (68) | 16.2 (80) | 956 | 4.8 (22) |
| Sauk Cty | 957 | 6.0 (10) | 9.9 (73) | 3,883 | 5.9 (72) | 2.2 (18) | 14.4 (72) | 1,681 | 18.3 (70) | 17.6 (85) | 2,759 | 5.9 (30) |
| Sawyer Cty | 294 | 6.8 (18) | 8.5 (48) | 1,293 | 4.5 (26) | 2.9 (34) | 11.0 (25) | 580 | 17.8 (64) | 13.3 (54) | 960 | 4.0 (16) |
| Seeds Of H | 2,412 | 7.4 (33) | 8.5 (48) | 9,507 | 3.2 (7) | 4.4 (79) | 11.4 (33) | 4,236 | 17.8 (64) | 14.5 (67) | 6,967 | 11.7 (77) |
| Shawano Ct | 558 | 8.4 (55) | 7.9 (35) | 2,296 | 5.3 (52) | 3.2 (52) | 12.8 (56) | 1,026 | 13.9 (16) | 10.7 (19) | 1,742 | 5.4 (25) |
| Sheboygan | 1,671 | 7.8 (40) | 6.8 (15) | 6,574 | 5.6 (64) | 5.5 (93) | 11.9 (44) | 2,937 | 16.0 (39) | 13.3 (54) | 4,818 | 3.9 (14) |
| St. Croix | 853 | 8.3 (52) | 9.6 (66) | 3,287 | 5.8 (70) | 2.8 (29) | 11.5 (34) | 1,391 | 17.2 (57) | 12.7 (44) | 2,299 | 3.8 (13) |
| SWCAP | 184 | 7.1 (22) | 6.0 (10) | 712 | 6.7 (92) | 2.8 (29) | 12.4 (48) | 287 | 24.4 (97) | 13.9 (60) | 496 | 10.5 (68) |
| Taylor Cty | 333 | 7.8 (40) | 9.9 (73) | 1,304 | 3.5 (9) | 6.1 (96) | 10.7 (21) | 583 | 11.0 (3) | 10.1 (15) | 972 | 3.1 (6) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | |
|---|-----------------|------------------------|------------------------|-------------------------|-------------------------|---------------------|----------------------|--------------------------------|----------------------------|-----------------------------|--------------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | High | | Short Stature (6) | Under- weight (7) | Obese (7) | | ≥ 2 Yrs Over- weight (7) | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- <95th (Rank) | % ≥95th (Rank) | Number % (Rank) |
| Trempealeau | 574 | 7.1 (22) | 9.9 (73) | 2,227 | 4.4 (22) | 4.6 (82) | 15.3 (83) | 984 | 19.5 (81) | 15.2 (74) | 1,583 10.0 (64) |
| Vernon Cou | 459 | 8.7 (66) | 13.5 (85) | 2,027 | 5.2 (49) | 2.8 (29) | 13.6 (65) | 916 | 17.1 (53) | 16.2 (80) | 1,473 5.6 (28) |
| Vilas Cty | 207 | 7.7 (38) | 7.7 (29) | 726 | 5.1 (44) | 3.6 (60) | 11.6 (35) | 279 | 15.4 (31) | 12.9 (46) | 459 2.8 (3) |
| Walworth C | 1,476 | 7.2 (27) | 8.6 (51) | 5,752 | 5.4 (57) | 2.7 (27) | 12.7 (55) | 2,350 | 15.6 (32) | 15.0 (72) | 3,978 7.1 (38) |
| Washburn C | 304 | 8.6 (62) | 8.9 (55) | 1,294 | 4.8 (34) | 6.3 (97) | 9.4 (8) | 591 | 11.2 (4) | 9.3 (12) | 943 5.3 (24) |
| Washington | 1,132 | 8.3 (52) | 6.7 (14) | 4,093 | 6.1 (80) | 3.7 (62) | 11.7 (39) | 1,738 | 16.3 (43) | 10.7 (19) | 2,875 8.5 (51) |
| Waukesha C | 1,968 | 8.2 (49) | 7.8 (32) | 7,109 | 5.1 (44) | 2.5 (23) | 13.4 (63) | 2,777 | 16.7 (47) | 13.2 (50) | 4,514 6.2 (31) |
| Waupaca Ct | 669 | 10.6 (81) | 8.4 (44) | 2,421 | 5.9 (72) | 3.8 (65) | 13.8 (68) | 1,000 | 13.2 (11) | 11.4 (28) | 1,643 4.0 (16) |
| Waushara | 338 | 8.0 (47) | 7.7 (29) | 1,377 | 5.1 (44) | 3.5 (58) | 14.6 (74) | 610 | 19.3 (80) | 17.9 (86) | 972 10.4 (66) |
| WC Capitol | 2,839 | 11.0 (83) | 5.2 (7) | 10,694 | 5.7 (68) | 4.2 (74) | 10.1 (15) | 4,405 | 12.9 (8) | 9.3 (12) | 7,295 19.9 (94) |
| WC North | 1,509 | 12.8 (86) | 4.1 (4) | 5,833 | 5.8 (70) | 4.3 (76) | 9.7 (9) | 2,484 | 13.5 (13) | 10.7 (19) | 4,177 25.0 (97) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

| Comparison of Growth and Anemia Indicators by Clinic (3,4) | | | | | | | | | | | |
|---|-----------------|------------------------|------------------------|-------------------------|-------------------------|---------------------|----------------------|--------------------------------|----------------------------|-----------------------------|---------------------|
| Clinic | Birthweight (5) | | | Height and Weight | | | | | | Anemia Low Hb/Hct (8) | |
| | Low | High | | Short Stature (6) | Under- weight (7) | Obese (7) | | ≥ 2 Yrs Over- weight (7) | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % ≥95th (Rank) | Number | % 85th- <95th (Rank) | % ≥95th (Rank) | Number % (Rank) |
| WC Teutonia | 1,663 | 13.3 (88) | 2.9 (1) | 6,744 | 7.8 (98) | 3.9 (68) | 12.6 (52) | 3,069 | 15.7 (35) | 11.7 (34) | 4,777 25.4 (98) |
| West Allis | 1,051 | 8.7 (66) | 8.7 (53) | 3,627 | 4.8 (34) | 4.8 (86) | 10.3 (17) | 1,410 | 16.5 (45) | 11.7 (34) | 2,382 13.0 (83) |
| Wood Cty | 1,199 | 7.1 (22) | 8.0 (38) | 4,216 | 5.6 (64) | 3.0 (40) | 11.3 (31) | 1,715 | 16.6 (46) | 11.1 (24) | 2,820 8.4 (49) |
| Unknown Clinics | 0 | * * | * * | 40 | * * | * * | * * | 21 | * * | * * | 33 * * |
| Wisconsin | 94,141 | 8.6 | 7.6 | 365,388 | 5.5 | 3.6 | 12.6 | 159,178 | 16.2 | 13.3 | 260,670 10.9 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic Children Aged < 5 Years ⁽³⁾ (2)

| Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic | | | | | | | | | | | | | | |
|---|----------------------|------------------------|--------|--|--------|---|--------------------------------|--|--------|--|-----------------------------------|-----------|-------------------------|-----------|
| | Breastfeeding (5) | | | | | | Exclusive Breastfeeding (6) | | | | ≤2 Hours TV Viewing/Day (7) | | Smoking in Household | |
| Clinic | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months | Number | % Breastfed At Least 3 Months | Number | % Breastfed At Least 6 Months | Number | % | Number | % |
| | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | |
| 16Th St Co | 4,656 | 84.5 (1) | 3,533 | 43.3 (1) | 3,645 | 26.3 (1) | 3,037 | 11.1 (50) | 3,161 | 5.6 (48) | 10,767 | 80.3 (68) | 10,518 | 6.3 (6) |
| Adams Cty | 378 | 64.3 (54) | 274 | 15.3 (78) | 261 | 7.3 (80) | 238 | 13.9 (34) | 254 | 4.7 (59) | 741 | 80.0 (71) | 807 | 23.9 (97) |
| Ashland Cty | 370 | 70.5 (21) | 281 | 30.6 (10) | 282 | 21.3 (3) | 243 | 21.0 (8) | 262 | 13.7 (2) | 842 | 88.1 (11) | 847 | 9.7 (34) |
| Aurora | 4,024 | 39.8 (89) | 2,086 | 11.9 (87) | 2,129 | 7.1 (82) | 1,854 | 3.5 (86) | 1,861 | 1.5 (84) | 4,531 | 83.0 (43) | 6,689 | 16.0 (81) |
| Barron Cty | 913 | 68.1 (37) | 604 | 22.8 (38) | 564 | 9.8 (59) | 569 | 15.5 (27) | 543 | 5.9 (44) | 1,471 | 80.8 (59) | 1,800 | 8.7 (28) |
| Bayfield Cty | 165 | 73.9 (11) | 116 | 38.8 (3) | 111 | 19.8 (4) | 103 | 33.0 (1) | 106 | 15.1 (1) | 381 | 85.3 (29) | 395 | 8.6 (26) |
| Beloit | 1,646 | 50.1 (84) | 1,077 | 19.3 (63) | 1,087 | 11.7 (39) | 929 | 4.7 (82) | 923 | 2.1 (80) | 2,652 | 85.3 (29) | 3,256 | 7.9 (20) |
| Buffalo Cty | 207 | 73.4 (14) | 138 | 21.7 (47) | 136 | 12.5 (29) | 126 | 10.3 (54) | 119 | 7.6 (23) | 380 | 87.6 (13) | 449 | 7.1 (12) |
| Burnett Cty | 288 | 76.7 (4) | 208 | 24.0 (33) | 204 | 12.3 (30) | 190 | 18.9 (15) | 185 | 9.2 (10) | 546 | 95.4 (2) | 626 | 14.2 (74) |
| Calumet Ct | 484 | 62.8 (56) | 315 | 22.2 (41) | 325 | 10.8 (48) | 289 | 14.5 (30) | 282 | 10.3 (7) | 922 | 80.8 (59) | 1,052 | 8.5 (25) |
| Chippewa C | 1,071 | 66.4 (44) | 703 | 20.6 (56) | 723 | 9.0 (66) | 602 | 13.1 (37) | 597 | 6.5 (36) | 1,991 | 87.6 (13) | 2,270 | 9.7 (34) |
| Clark Cty | 585 | 73.3 (15) | 395 | 19.2 (64) | 391 | 7.2 (81) | 366 | 12.8 (41) | 371 | 3.2 (69) | 1,137 | 84.4 (34) | 1,240 | 9.8 (38) |
| Columbia C | 714 | 69.6 (30) | 476 | 26.7 (19) | 439 | 13.9 (19) | 407 | 16.7 (23) | 391 | 6.9 (33) | 1,109 | 81.7 (53) | 1,488 | 10.3 (50) |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 2
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (4,5) | | | | | | | | | | | | | | |
|--|----------------------|------------------------|--------|--|--------|---|--------------------------------|--|--------|--|-----------------------------------|-----------|-------------------------|-----------|
| | Breastfeeding (5) | | | | | | Exclusive Breastfeeding (6) | | | | ≤2 Hours TV Viewing/Day (7) | | Smoking in Household | |
| Clinic | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months | Number | % Breastfed At Least 3 Months | Number | % Breastfed At Least 6 Months | Number | % | Number | % |
| | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) |
| Crawford Cty | 265 | 64.9 (50) | 162 | 22.2 (41) | 166 | 16.3 (8) | 145 | 17.2 (20) | 143 | 7.0 (29) | 417 | 85.6 (26) | 556 | 8.6 (26) |
| Cudahy | 1,338 | 65.6 (48) | 824 | 20.0 (59) | 829 | 9.4 (63) | 653 | 8.0 (67) | 641 | 3.9 (66) | 1,805 | 70.5 (95) | 2,551 | 23.3 (96) |
| Dane Cty | 633 | 73.5 (13) | 566 | 32.3 (6) | 632 | 18.5 (5) | 410 | 11.5 (45) | 443 | 7.0 (29) | 1,713 | 79.3 (75) | 2,233 | 7.8 (19) |
| Dodge Cty | 1,056 | 74.5 (10) | 673 | 28.1 (15) | 641 | 16.1 (10) | 642 | 20.9 (9) | 616 | 11.0 (5) | 1,748 | 88.4 (10) | 2,214 | 7.1 (12) |
| Door Cty | 347 | 71.2 (19) | 216 | 29.6 (11) | 237 | 11.4 (42) | 206 | 21.4 (3) | 200 | 8.0 (18) | 680 | 84.3 (35) | 770 | 7.9 (20) |
| Douglas Cty | 910 | 69.5 (31) | 617 | 26.9 (17) | 616 | 13.6 (23) | 597 | 21.4 (3) | 576 | 13.7 (2) | 1,696 | 88.6 (8) | 1,975 | 10.1 (45) |
| Dunn Cty | 690 | 68.6 (34) | 478 | 22.8 (38) | 470 | 9.8 (59) | 444 | 20.5 (10) | 434 | 11.1 (4) | 1,389 | 80.7 (61) | 1,577 | 6.2 (5) |
| Eau Claire | 1,793 | 70.6 (20) | 1,213 | 20.9 (51) | 1,240 | 8.8 (69) | 1,104 | 11.2 (48) | 1,099 | 5.4 (51) | 3,329 | 84.3 (35) | 3,763 | 7.0 (10) |
| Florence C | 79 | * * | 54 | * * | 51 | * * | 43 | * * | 43 | * * | 193 | 88.6 (8) | 205 | 10.2 (49) |
| Fond Du La | 1,513 | 67.7 (40) | 1,033 | 24.3 (26) | 1,033 | 11.2 (45) | 995 | 16.8 (21) | 961 | 6.3 (37) | 2,785 | 86.0 (22) | 3,205 | 10.1 (45) |
| Forest Cty | 165 | 53.3 (79) | 119 | 20.2 (57) | 121 | 7.4 (79) | 101 | 11.9 (42) | 104 | 4.8 (57) | 388 | 76.0 (88) | 379 | 19.3 (91) |
| GLITC-Bad | 74 | * * | 55 | * * | 59 | * * | 41 | * * | 43 | * * | 204 | 79.4 (74) | 183 | 18.0 (86) |
| GLITC-Bara | 45 | * * | 30 | * * | 24 | * * | 22 | * * | 20 | * * | 62 | * * | 87 | * * |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 - 2008 Pediatric Nutrition Surveillance Wisconsin

Page: 3
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | Exclusive Breastfeeding (6) | | ≤2 Hours TV Viewing/Day (7) | Smoking in Household |
|------------|--|--|---|--|--|-----------------------------------|-------------------------------|
| Clinic | Number % Ever Breastfed (Rank) | Number % Breastfed At Least 6 Months (Rank) | Number % Breastfed At Least 12 Months (Rank) | Number % Breastfed At Least 3 Months (Rank) | Number % Breastfed At Least 6 Months (Rank) | Number % (Rank) | Number % (Rank) |
| GLITC-BRF | 81 * * | 67 * * | 69 * * | 55 * * | 58 * * | 197 80.7 (61) | 197 17.8 (84) |
| GLITC-CEX | 25 * * | 17 * * | 16 * * | 17 * * | 16 * * | 45 * * | 51 * * |
| GLITC-LCO | 139 48.9 (87) | 94 * * | 109 3.7 (89) | 80 * * | 85 * * | 220 98.6 (1) | 270 11.9 (62) |
| GLITC-Mole | 23 * * | 19 * * | 19 * * | 16 * * | 17 * * | 61 * * | 56 * * |
| GLITC-Pete | 192 56.3 (75) | 154 17.5 (74) | 153 13.1 (25) | 133 7.5 (70) | 137 4.4 (60) | 389 77.4 (79) | 392 18.9 (89) |
| GLITC-Pota | 59 * * | 38 * * | 39 * * | 33 * * | 32 * * | 121 81.8 (52) | 129 22.5 (94) |
| GLITC-Red | 79 * * | 58 * * | 69 * * | 43 * * | 41 * * | 189 82.0 (50) | 178 14.0 (71) |
| GLITC-St.C | 93 * * | 74 * * | 80 * * | 67 * * | 66 * * | 65 * * | 242 13.6 (65) |
| GLITC-Stoc | 91 * * | 63 * * | 63 * * | 54 * * | 56 * * | 215 77.2 (82) | 209 12.0 (63) |
| Grant Cty | 867 62.6 (57) | 617 23.8 (34) | 592 11.8 (37) | 423 13.2 (36) | 427 7.0 (29) | 1,720 82.7 (46) | 1,912 9.0 (29) |
| Green Cty | 523 60.2 (64) | 359 21.2 (50) | 375 12.0 (35) | 337 17.8 (17) | 335 6.3 (37) | 895 87.5 (15) | 1,099 15.7 (79) |
| Green Lake | 261 52.9 (80) | 173 13.3 (83) | 165 6.7 (84) | 116 6.0 (75) | 117 4.3 (61) | 419 75.2 (91) | 526 9.7 (34) |
| Greenfield | 788 65.9 (45) | 431 24.1 (30) | 412 10.7 (49) | 378 10.1 (56) | 361 5.3 (53) | 987 75.8 (90) | 1,425 17.1 (83) |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 - 2008 Pediatric Nutrition Surveillance (1)
Wisconsin

Page: 4
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | | | | Exclusive Breastfeeding (6) | | | | ≤2 Hours TV Viewing/Day (7) | | Smoking in Household | |
|------------|----------------------|------------------------|--------|--|--------|---|--------------------------------|--|--------|--|-----------------------------------|-----------|-------------------------|-----------|
| Clinic | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months | Number | % Breastfed At Least 3 Months | Number | % Breastfed At Least 6 Months | Number | % | Number | % |
| | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | |
| Hlth & Nut | 3,046 | 61.0 (62) | 1,879 | 18.6 (66) | 1,805 | 10.5 (52) | 1,741 | 8.0 (67) | 1,707 | 3.7 (67) | 4,577 | 77.3 (81) | 5,776 | 11.0 (55) |
| Iowa Cty | 328 | 65.9 (45) | 208 | 22.6 (40) | 183 | 12.6 (28) | 194 | 15.5 (27) | 191 | 7.9 (21) | 409 | 86.6 (17) | 621 | 9.8 (38) |
| Iron Cty | 73 | * * | 50 | * * | 44 | * * | 49 | * * | 41 | * * | 146 | 92.5 (3) | 171 | 9.4 (31) |
| Jackson Ct | 406 | 70.4 (23) | 289 | 22.1 (43) | 290 | 11.4 (42) | 248 | 11.3 (47) | 243 | 5.8 (47) | 842 | 85.7 (25) | 887 | 11.0 (55) |
| Janesville | 1,741 | 52.9 (80) | 1,039 | 17.7 (72) | 1,035 | 9.7 (61) | 928 | 8.6 (64) | 872 | 4.2 (62) | 2,464 | 85.0 (32) | 3,305 | 13.8 (68) |
| Jefferson | 1,355 | 69.4 (33) | 984 | 24.9 (24) | 908 | 13.8 (21) | 830 | 7.2 (71) | 826 | 5.1 (54) | 2,394 | 86.5 (18) | 2,881 | 9.8 (38) |
| Juneau | 578 | 69.9 (28) | 427 | 24.1 (30) | 394 | 10.9 (47) | 402 | 16.4 (24) | 398 | 8.5 (14) | 1,103 | 82.8 (45) | 1,258 | 20.0 (92) |
| Kenosha | 3,419 | 59.5 (67) | 2,039 | 18.5 (67) | 2,046 | 8.9 (68) | 1,956 | 5.6 (77) | 1,848 | 2.9 (75) | 5,213 | 81.1 (57) | 6,892 | 13.6 (65) |
| Kewaunee C | 237 | 71.3 (18) | 152 | 24.3 (26) | 146 | 12.3 (30) | 122 | 21.3 (6) | 120 | 8.3 (17) | 467 | 69.2 (96) | 521 | 9.4 (31) |
| La Crosse | 1,894 | 70.1 (26) | 1,233 | 24.6 (25) | 1,185 | 13.1 (25) | 1,025 | 11.2 (48) | 988 | 5.0 (56) | 3,293 | 86.4 (20) | 3,927 | 4.3 (1) |
| Langlade | 377 | 54.6 (76) | 262 | 13.0 (84) | 264 | 7.6 (77) | 219 | 7.8 (69) | 222 | 3.2 (69) | 667 | 81.1 (57) | 808 | 16.6 (82) |
| Lincoln | 372 | 61.3 (60) | 232 | 18.1 (68) | 204 | 8.8 (69) | 201 | 10.0 (57) | 192 | 4.2 (62) | 636 | 86.2 (21) | 761 | 15.0 (76) |
| Madison WI | 5,377 | 75.8 (7) | 3,342 | 39.3 (2) | 3,125 | 23.4 (2) | 2,993 | 14.2 (32) | 2,826 | 9.8 (9) | 7,492 | 85.4 (28) | 9,556 | 7.1 (12) |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 - 2008 Pediatric Nutrition Surveillance Wisconsin

Page: 5
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | | | | Exclusive Breastfeeding (6) | | | | ≤2 Hours TV Viewing/Day (7) | | Smoking in Household | |
|-------------|----------------------|------------------------|--------|--|--------|---|--------------------------------|--|--------|--|-----------------------------------|-----------|-------------------------|-----------|
| Clinic | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months | Number | % Breastfed At Least 3 Months | Number | % Breastfed At Least 6 Months | Number | % | Number | % |
| | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | |
| Manitowoc | 1,201 | 58.7 (70) | 769 | 19.6 (60) | 812 | 11.7 (39) | 723 | 11.1 (50) | 708 | 5.9 (44) | 2,259 | 77.0 (86) | 2,550 | 12.7 (64) |
| Marathon | 2,051 | 57.6 (72) | 1,426 | 17.0 (75) | 1,290 | 9.0 (66) | 1,251 | 9.3 (59) | 1,206 | 4.1 (64) | 3,512 | 80.5 (66) | 4,240 | 8.1 (22) |
| Marinette | 735 | 49.3 (86) | 519 | 16.2 (77) | 554 | 6.9 (83) | 462 | 9.5 (58) | 476 | 5.5 (50) | 1,542 | 86.0 (22) | 1,711 | 15.2 (77) |
| Marquette | 182 | 59.9 (66) | 106 | 20.8 (52) | 114 | 14.9 (15) | 81 | * * | 65 | * * | 324 | 76.5 (87) | 387 | 15.8 (80) |
| Menominee | 434 | 67.3 (43) | 301 | 21.3 (49) | 321 | 10.3 (54) | 256 | 2.7 (87) | 275 | 1.5 (84) | 836 | 90.2 (6) | 861 | 10.7 (52) |
| MilwaukeeJ | 1,738 | 69.9 (28) | 1,204 | 25.9 (21) | 1,261 | 14.4 (16) | 1,050 | 6.8 (72) | 1,043 | 3.0 (73) | 3,723 | 77.2 (82) | 3,786 | 11.4 (59) |
| MilwaukeeK | 1,701 | 44.9 (88) | 1,188 | 9.8 (88) | 1,283 | 5.5 (88) | 1,022 | 3.7 (84) | 1,051 | 1.0 (86) | 3,527 | 64.3 (97) | 3,801 | 21.0 (93) |
| MilwaukeeN | 3,116 | 50.1 (84) | 1,935 | 12.0 (86) | 1,972 | 8.0 (75) | 1,795 | 3.6 (85) | 1,732 | 1.8 (82) | 4,719 | 72.3 (94) | 6,183 | 13.9 (69) |
| MilwaukMLK | 2,322 | 56.5 (73) | 1,615 | 14.1 (80) | 1,650 | 8.2 (73) | 1,419 | 4.9 (80) | 1,396 | 2.1 (80) | 4,144 | 74.5 (92) | 4,838 | 23.1 (95) |
| Monroe Cou | 897 | 68.2 (35) | 578 | 20.1 (58) | 570 | 11.2 (45) | 527 | 10.4 (53) | 498 | 5.4 (51) | 1,399 | 82.3 (49) | 1,735 | 11.4 (59) |
| N.E.W. East | 2,204 | 64.9 (50) | 1,516 | 21.9 (44) | 1,642 | 13.9 (19) | 1,380 | 5.7 (76) | 1,357 | 3.2 (69) | 4,487 | 77.2 (82) | 4,766 | 5.9 (4) |
| N.E.W. West | 2,437 | 58.1 (71) | 1,570 | 19.6 (60) | 1,543 | 10.5 (52) | 1,337 | 8.8 (63) | 1,328 | 3.6 (68) | 4,066 | 79.8 (73) | 4,858 | 11.0 (55) |
| Neenah | 1,038 | 67.4 (42) | 625 | 21.8 (46) | 569 | 9.3 (65) | 607 | 12.9 (40) | 588 | 6.8 (35) | 1,565 | 82.6 (47) | 2,040 | 13.9 (69) |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 - 2008 Pediatric Nutrition Surveillance Wisconsin

Page: 6
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | | | | Exclusive Breastfeeding (6) | | | | ≤2 Hours TV Viewing/Day (7) | | Smoking in Household | |
|------------|----------------------|------------------------|--------|--|--------|---|--------------------------------|--|--------|--|-----------------------------------|-----------|-------------------------|-----------|
| Clinic | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months | Number | % Breastfed At Least 3 Months | Number | % Breastfed At Least 6 Months | Number | % | Number | % |
| | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | |
| Oconto Cty | 464 | 53.4 (78) | 331 | 13.0 (84) | 308 | 5.8 (86) | 308 | 5.2 (79) | 310 | 2.3 (79) | 840 | 81.5 (54) | 999 | 14.0 (71) |
| Oneida Cty | 499 | 60.9 (63) | 313 | 23.3 (36) | 289 | 11.8 (37) | 305 | 20.0 (12) | 289 | 8.0 (18) | 768 | 92.3 (4) | 1,034 | 9.8 (38) |
| Oneida Tri | 454 | 62.6 (57) | 297 | 17.8 (71) | 327 | 7.6 (77) | 272 | 11.4 (46) | 262 | 6.1 (41) | 758 | 80.3 (68) | 910 | 17.9 (85) |
| Oshkosh | 1,127 | 59.0 (69) | 734 | 16.9 (76) | 702 | 8.4 (72) | 650 | 8.6 (64) | 665 | 2.9 (75) | 1,743 | 83.9 (40) | 2,272 | 9.0 (29) |
| Outagamie | 2,095 | 64.8 (52) | 1,290 | 20.8 (52) | 1,321 | 10.3 (54) | 1,267 | 14.4 (31) | 1,205 | 6.0 (43) | 3,648 | 80.5 (66) | 4,419 | 7.4 (17) |
| Ozaukee | 458 | 75.8 (7) | 275 | 31.3 (7) | 262 | 14.1 (17) | 280 | 24.6 (2) | 264 | 7.6 (23) | 640 | 80.3 (68) | 879 | 7.2 (15) |
| Pepin | 89 | * * | 57 | * * | 56 | * * | 43 | * * | 45 | * * | 190 | 86.8 (16) | 195 | 9.7 (34) |
| Pierce Cty | 514 | 67.9 (39) | 357 | 21.6 (48) | 347 | 8.1 (74) | 350 | 17.7 (19) | 324 | 8.0 (18) | 1,045 | 90.0 (7) | 1,179 | 10.9 (54) |
| Polk Cty | 776 | 75.9 (6) | 510 | 25.9 (21) | 466 | 10.7 (49) | 444 | 21.4 (3) | 436 | 8.5 (14) | 1,154 | 79.3 (75) | 1,516 | 10.0 (44) |
| Portage Ct | 937 | 68.2 (35) | 619 | 24.1 (30) | 555 | 13.0 (27) | 576 | 13.5 (35) | 540 | 6.1 (41) | 1,595 | 88.1 (11) | 1,926 | 5.5 (2) |
| Price Cty | 282 | 70.2 (24) | 233 | 31.3 (7) | 233 | 15.9 (11) | 172 | 11.6 (44) | 180 | 7.2 (28) | 745 | 79.9 (72) | 703 | 13.7 (67) |
| Racine Hea | 1,317 | 52.8 (82) | 863 | 17.7 (72) | 813 | 10.0 (56) | 770 | 4.9 (80) | 761 | 2.8 (77) | 2,351 | 77.8 (77) | 2,714 | 10.1 (45) |
| Richland | 290 | 77.6 (2) | 181 | 24.3 (26) | 171 | 14.0 (18) | 163 | 17.8 (17) | 162 | 9.9 (8) | 479 | 81.2 (56) | 587 | 6.8 (9) |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 - 2008 Pediatric Nutrition Surveillance Wisconsin

Page: 7
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | | | | Exclusive Breastfeeding (6) | | | | ≤2 Hours TV Viewing/Day (7) | | Smoking in Household | |
|-------------|----------------------|------------------------|--------|--|--------|---|--------------------------------|--|--------|--|-----------------------------------|-----------|-------------------------|-----------|
| Clinic | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months | Number | % Breastfed At Least 3 Months | Number | % Breastfed At Least 6 Months | Number | % | Number | % |
| | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | |
| Rusk Cty | 301 | 61.1 (61) | 194 | 19.6 (60) | 199 | 11.6 (41) | 157 | 10.8 (52) | 159 | 6.3 (37) | 661 | 85.6 (26) | 702 | 14.1 (73) |
| Sauk Cty | 1,078 | 76.1 (5) | 757 | 32.4 (5) | 669 | 15.5 (13) | 758 | 21.2 (7) | 694 | 8.9 (12) | 1,806 | 84.3 (35) | 2,195 | 27.8 (98) |
| Sawyer Cty | 322 | 72.0 (16) | 229 | 19.2 (64) | 222 | 9.9 (57) | 210 | 16.2 (25) | 201 | 8.5 (14) | 631 | 80.7 (61) | 689 | 11.6 (61) |
| Seeds Of H | 2,549 | 71.4 (17) | 1,760 | 23.8 (34) | 1,886 | 12.2 (32) | 1,591 | 6.2 (74) | 1,610 | 3.0 (73) | 4,611 | 82.9 (44) | 5,060 | 8.1 (22) |
| Shawano Ct | 606 | 65.8 (47) | 415 | 20.7 (55) | 432 | 10.6 (51) | 364 | 9.3 (59) | 373 | 7.0 (29) | 1,132 | 85.2 (31) | 1,300 | 10.8 (53) |
| Sheboygan | 1,823 | 64.5 (53) | 1,240 | 24.3 (26) | 1,299 | 13.5 (24) | 1,146 | 9.3 (59) | 1,113 | 4.8 (57) | 3,155 | 80.7 (61) | 3,712 | 9.9 (43) |
| St. Croix | 933 | 73.6 (12) | 588 | 28.9 (13) | 554 | 15.7 (12) | 562 | 18.3 (16) | 527 | 8.7 (13) | 1,476 | 77.4 (79) | 1,932 | 7.0 (10) |
| SWCAP | 186 | 65.1 (49) | 129 | 26.4 (20) | 123 | 12.2 (32) | 117 | 10.3 (54) | 119 | 5.9 (44) | 319 | 84.3 (35) | 416 | 8.4 (24) |
| Taylor Cty | 348 | 68.1 (37) | 219 | 21.9 (44) | 225 | 13.8 (21) | 186 | 15.6 (26) | 184 | 7.6 (23) | 623 | 80.7 (61) | 756 | 18.8 (88) |
| Trempealeau | 598 | 70.2 (24) | 388 | 28.9 (13) | 373 | 16.4 (7) | 338 | 11.8 (43) | 328 | 4.0 (65) | 1,075 | 91.8 (5) | 1,292 | 5.7 (3) |
| Vernon Cou | 510 | 74.7 (9) | 359 | 34.0 (4) | 357 | 16.5 (6) | 315 | 20.3 (11) | 299 | 10.7 (6) | 1,002 | 84.9 (33) | 1,107 | 7.2 (15) |
| Vilas Cty | 221 | 54.3 (77) | 140 | 22.9 (37) | 127 | 9.4 (63) | 137 | 16.8 (21) | 130 | 9.2 (10) | 316 | 83.5 (41) | 405 | 10.1 (45) |
| Walworth C | 1,659 | 77.1 (3) | 1,131 | 31.3 (7) | 1,006 | 15.2 (14) | 1,082 | 19.5 (13) | 1,050 | 5.6 (48) | 2,544 | 83.1 (42) | 3,363 | 6.5 (8) |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2006 - 2008 Pediatric Nutrition Surveillance Wisconsin

Page: 8
Run Date: 3 / 13 / 09

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
(4,5)

| | Breastfeeding (5) | | | | | | Exclusive Breastfeeding (6) | | | | ≤2 Hours TV Viewing/Day (7) | | Smoking in Household | |
|-----------------|----------------------|------------------------|--------|--|--------|---|--------------------------------|--|--------|--|-----------------------------------|-----------|-------------------------|-----------|
| Clinic | Number | % Ever Breastfed | Number | % Breastfed At Least 6 Months | Number | % Breastfed At Least 12 Months | Number | % Breastfed At Least 3 Months | Number | % Breastfed At Least 6 Months | Number | % | Number | % |
| | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) | | (Rank) |
| Washburn C | 342 | 70.5 (21) | 256 | 27.0 (16) | 252 | 9.9 (57) | 211 | 14.2 (32) | 215 | 5.1 (54) | 635 | 77.8 (77) | 720 | 6.3 (6) |
| Washington | 1,253 | 70.0 (27) | 740 | 26.9 (17) | 768 | 12.0 (35) | 723 | 19.1 (14) | 696 | 7.6 (23) | 1,864 | 86.5 (18) | 2,415 | 9.4 (31) |
| Waukesha C | 2,174 | 69.5 (31) | 1,371 | 29.0 (12) | 1,191 | 16.3 (8) | 1,296 | 13.0 (38) | 1,266 | 6.9 (33) | 2,994 | 81.9 (51) | 4,276 | 7.5 (18) |
| Waupaca Ct | 713 | 61.7 (59) | 443 | 20.8 (52) | 421 | 9.7 (61) | 401 | 13.0 (38) | 390 | 6.2 (40) | 1,067 | 85.8 (24) | 1,402 | 15.5 (78) |
| Waushara | 372 | 59.4 (68) | 238 | 18.1 (68) | 222 | 8.6 (71) | 176 | 9.1 (62) | 164 | 7.3 (27) | 648 | 73.9 (93) | 780 | 10.5 (51) |
| WC Capitol | 3,060 | 60.0 (65) | 1,932 | 14.9 (79) | 2,007 | 7.8 (76) | 1,676 | 6.4 (73) | 1,705 | 2.5 (78) | 4,779 | 77.1 (85) | 6,011 | 11.2 (58) |
| WC North | 1,618 | 51.5 (83) | 1,062 | 13.7 (81) | 1,106 | 6.2 (85) | 900 | 4.1 (83) | 928 | 0.8 (87) | 2,685 | 84.3 (35) | 3,155 | 19.0 (90) |
| WC Teutonia | 1,804 | 64.3 (54) | 1,240 | 13.5 (82) | 1,257 | 5.7 (87) | 1,081 | 5.3 (78) | 1,074 | 1.7 (83) | 3,315 | 76.0 (88) | 3,701 | 18.5 (87) |
| West Allis | 1,162 | 67.6 (41) | 653 | 25.3 (23) | 647 | 12.1 (34) | 655 | 8.1 (66) | 592 | 3.2 (69) | 1,508 | 81.4 (55) | 2,144 | 14.7 (75) |
| Wood Cty | 1,282 | 56.5 (73) | 741 | 18.1 (68) | 729 | 11.4 (42) | 670 | 14.9 (29) | 627 | 7.8 (22) | 1,861 | 82.4 (48) | 2,469 | 9.8 (38) |
| Unknown Clinics | 0 | * * | 2 | * * | 2 | * * | 0 | * * | 2 | * * | 21 | * * | 30 | * * |
| Wisconsin | 101,105 | 64.2 | 66,185 | 22.9 | 65,711 | 12.2 | 59,278 | 10.7 | 58,212 | 5.1 | 172,041 | 81.3 | 206,306 | 11.3 |

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.